

Monitoring the Sustainable Intensification of Arable Agriculture: the Potential Role of Earth Observation

Merryn L. Hunt^{*a,b}, George Alan Blackburn^b, Clare S. Rowland^a

^a Centre for Ecology and Hydrology, Lancaster Environment Centre, Library Avenue, Bailrigg, Lancaster, United Kingdom, LA1 4AP

^b Lancaster Environment Centre, Library Avenue, Lancaster University, Lancaster, United Kingdom, LA1 4YQ

*Corresponding author email address: m.hunt3@lancaster.ac.uk (M. L. Hunt)

Abstract

Sustainable intensification (SI) has been proposed as a possible solution to the conflicting problems of meeting projected increase in food demand and preserving environmental quality. SI would provide necessary production increases while simultaneously reducing or eliminating environmental degradation, without taking land from competing demands. An important component of achieving these aims is the development of suitable methods for assessing the temporal variability of both the intensification and sustainability of agriculture. Current assessments rely on traditional data collection methods that produce data of limited spatial and temporal resolution. Earth Observation (EO) provides a readily accessible, long-term dataset with global coverage at various spatial and temporal resolutions. In this paper we demonstrate how EO could significantly contribute to SI assessments, providing opportunities to quantify agricultural intensity and environmental sustainability. We review an extensive body of research on EO-based methods to assess multiple indicators of both agricultural intensity and environmental sustainability. To date these techniques have not been combined to assess SI; here we identify the opportunities and initial steps required to achieve this. In this context, we propose a development of a set of essential sustainable intensification variables (ESIVs) that could be derived from EO data.

Keywords: Sustainable intensification assessment; Earth Observation; agricultural intensity indicators; environmental sustainability indicators; crop yield; vegetation health; soil quality; water quality; biodiversity; ecosystem health

1. Introduction

With a projected population increase of 2.3 billion by 2050, increases in income and shifting food consumption patterns, global food production must increase by an estimated 70% to meet growing demand (Campbell *et al.*, 2014; Caviglia and Andrade, 2010; Dillon *et al.*, 2016; FAO, 2009; Garnett *et al.*, 2013; Lampkin *et al.*, 2015; Schut *et al.*, 2016; Tilman *et al.*, 2011, 2002). Both agricultural expansion (clearing additional land for crop production) and intensification (increasing productivity of existing agricultural land) could provide necessary crop production increases (Godfray and Garnett, 2014; Tilman *et al.*, 2011). Current competition for land restricts the potential for conversion of new land to agriculture, thus limiting the viability of expansion in many cases (Godfray *et al.*, 2010; Pretty *et al.*, 2011). In addition, expansion is thought to have a greater detrimental impact on the environment than intensification, with potential for significant greenhouse gas release through land conversion and major biodiversity losses affecting essential ecosystem service provision (Garnett *et al.*, 2013; Godfray and Garnett, 2014). Consequently, future demands must be met through production increases on current agricultural land alongside shifts in diet and the reduction in food waste (including transport and consumption). Previous agricultural intensification (AI) has been achieved through changes in management practices including increased agrochemical

inputs, cropping intensity and irrigation, and adoption of monoculture practices (Benton *et al.*, 2003; Crowder and Jabbour, 2014; Meeus, 1993; Stoate *et al.*, 2001). However, it is now widely recognised that such intensification measures detrimentally impact the environment, through over exploitation of natural resources for inputs and emission of pollution and waste (Pretty *et al.*, 2011). This raises concerns over the long-term ability to maintain intensive agricultural practices, with intensification-induced environmental degradation having negative feedbacks on sustained crop productivity (Bommarco *et al.*, 2013; Foley *et al.*, 2005; Matson *et al.*, 1997). It is clear therefore that a sustainable method of agricultural intensification is required.

One possible solution is sustainable intensification (SI), which involves increasing production efficiency to achieve higher agricultural outputs with the same or fewer inputs, while simultaneously significantly reducing or eliminating environmental degradation (Dillon *et al.*, 2016). However, no definitive mechanisms of SI exist, with the success of different methods dependent on situation-specific conditions. As such, to ensure any attempts at SI are successful suitable methods are required to assess the sustainability of intensification efficiently over diverse landscapes and spatial scales on a long-term basis.

The purpose of this review is to outline the current state of SI assessments and explore the potential contributions EO could make. While it is true that various studies have used EO to assess either agricultural intensity or various indicators of environmental sustainability, to date no one has attempted to combine established EO-based methods to provide an actual assessment of sustainable intensification. Hence, this review explores the basis for the development of an operational SI monitoring system that uses EO data. This review is structured as follows. Section 2 provides an overview of the key concepts of agricultural intensification (AI) and SI, as well as briefly introducing SI assessment. Section 3 and Section 4 present a more detailed outline of the current approaches used to assess agricultural intensity and agricultural environmental sustainability, respectively, highlighting ways in which EO data is presently used and further contributions it could make. The review concludes with a discussion of the opportunities of EO to contribute towards an operational SI monitoring system applicable for a range of spatial and temporal scales. This review focuses on the intensification arable agriculture; as such methods for monitoring pastoral agriculture are not explicitly discussed.

2. Key concepts/definitions

2.1 Agricultural Intensification (AI)

Agricultural intensification (AI) is the “increase in agricultural production per unit of inputs”, where inputs may include labour, land, time, seed, fertiliser, feed or cash (FAO, 2004). Intensification can refer to maintenance of current production with decreased inputs, and/or increased production through higher input productivity (FAO, 2004). Methods of intensification include: increased agrochemical inputs; increased cropping intensity (e.g. double or triple cropping); increased crop density; removal of linear and point features such as hedgerows and ponds (landscape simplification and field enlargement); decreased crop diversity (monoculture adoption); and increased irrigation (e.g. Crowder and Jabbour, 2014; Donald *et al.*, 2001; Newton, 2004; Stoate *et al.*, 2001). Detrimental environmental impacts of AI are wide-ranging, covering a range of spatial (local to global) and temporal (short-term to long-term or permanent) scales. These factors are driving the growing interest in alternative, more sustainable methods for meeting growing food demand. Figure S1 in the supplementary material highlights some of the key environmental impacts of various mechanisms of agricultural intensification. The potential for environmental degradation resulting from AI activities is intensified by the complexity of the agricultural environment, with numerous interactions, connections and feedbacks within the system, and multiple causal relationships. Such degradation could have significant impacts both within the immediate vicinity of intensification and over wider spatial scales. The wide range of potential impacts and system complexity poses a challenge when trying to devise a monitoring system that can accurately measure all required elements.

2.2 Sustainable Intensification (SI)

Sustainable Intensification (SI) has been proposed as an alternative to conventional intensification providing necessary yield increases, whilst ensuring environmental degradation is kept at a sustainable level (Tilman *et al.*, 2011). The concept originated in the 1990s (Buckwell *et al.*, 2014), with much debate since over the exact definition of the term “sustainable intensification”. A common definition describes SI as a form of production wherein greater yields are produced with the same or fewer inputs, while adverse environmental impacts are simultaneously reduced or eliminated and contribution to natural capital and ecosystem service flow is increased (Barnes and Thomson, 2014; Dillon *et al.*, 2016; Garnett and Godfray, 2012; Godfray *et al.*, 2010; Pretty, 2008; Pretty *et al.*, 2011). SI as a concept prescribes no particular development paths or methods; the aim is simply to create resource efficient agriculture with significantly better environmental performance than conventional intensification (Buckwell *et al.*, 2014). Instead, a framework is provided facilitating exploration of the optimum mix of approaches based on existing situation-specific biophysical, social, cultural and economic contexts (Buckwell *et al.*, 2014; Garnett *et al.*, 2013; Garnett and Godfray, 2012; Pretty and Bharucha, 2014). The suitability of different methods varies depending on conditions, as well as current agricultural productivity and environmental performance of the system (Buckwell *et al.*, 2014; Garnett and Godfray, 2012).

Possible interventions to achieve SI include: Integrated Pest Management (IPM), use of on- and off-farm biodiversity to manipulate pest ecologies; Agroforestry Systems, for example intercropping; Precision Agriculture; and Conservation Agriculture (Pretty and Bharucha, 2014).

2.3 Assessing Sustainable Intensification

As there is no definitive mechanism for SI, realising the goal of resource efficient agriculture requires suitable methods to assess the sustainability of intensification efficiently over diverse landscapes and spatial scales on a long-term basis. Current assessment attempts rely largely on farm surveys (questionnaires and interviews), field data, national government statistics and other traditional data sources. Data collection is often costly and time consuming, limiting the spatial and temporal scale and extent, with consequent impacts on the representativeness of both the data and the assessments. As such, the current reliance on interpolation of point data and average statistics severely restricts timely provision of accurate sustainability assessments for all agricultural areas. Deficiencies such as these highlight the need for a new, more efficient assessment technique.

Generally, studies focus on either agricultural sustainability, with no explicit attempt to quantify intensification (e.g. Dillon *et al.*, 2010; Rasul and Thapa, 2004), or on agricultural intensity at a specific point in time, with no assessment of sustainability (e.g. Herzog *et al.*, 2006; Niedertscheider *et al.*, 2016). Studies that assess the sustainability of intensification are largely conducted on farms where management practices are known to have shifted towards more intensive measures; hence, no attempt is made to quantify the degree or rate of intensification. The few studies that do include a measure of intensity (e.g. crop yield) commonly ignore change over time and use a single point in time. Intensification is a process rather than a fixed end state; looking at intensity at one specific time is therefore not sufficient (Elliott *et al.*, 2013; Firbank *et al.*, 2013). To get a full picture of the sustainability of any intensification in agricultural production, the change that has occurred over time must be investigated and trends identified to determine the actual environmental impacts. The potential of a quantified SI assessment approach at the individual farm level, using data held by farmers from two different years, was explored by Firbank *et al.* (2013). Results demonstrated the ability to assess sustainability of farms adopting different management strategies, but the data source restricted the completeness of the assessment, limiting measurable indicators and identifiable spatial and temporal variation. The limited temporal resolution of the data restricted full exploration of the intensification and subsequent effects on yield and environmental sustainability. The study constitutes an important step in the development of an operational SI assessment method, but more comprehensive data is required.

Ideally, field measurements would be taken at all locations to provide data on which SI could be assessed. However, as this is not feasible, data sources with greater spatial and temporal coverage and lower acquisition costs must be sought that can be used in conjunction with field measurements. Incorporation of satellite data into sustainability assessments could allow greater flexibility, over various spatial and temporal scales, providing more accurate and representative results at lower costs. Recent decades have seen considerable increases in Earth Observation (EO) data use, with applications in diverse research areas. A number of international monitoring systems have been developed incorporating satellite data for crop condition monitoring and yield forecasting over regional, national and global scales. Such systems include the Group on Earth Observations Global Agricultural Monitoring system (GEOGLAM) (GEO, n.d.), the USDA Foreign Agricultural Service (FAS) Global Agricultural Monitoring (GLAM) System (USDA FAS, n.d.), the Chinese Academy of Sciences Crop Watch Program (Bingfang, 2006), and the Monitoring Agriculture by Remote Sensing (MARS) project developed by the Joint Research Centre of the European Commission (European Commission, 2016). The operational status of these systems demonstrates the value of EO for agricultural monitoring. Outputs from these systems could be used to monitor intensification, but currently this is not done explicitly. The potential contribution of these EO-based systems to SI assessment has so far not been fully explored or realised, with little or no evidence of the use of satellite-derived data within agricultural sustainability assessments.

To date EO researchers have not explicitly attempted to quantify SI and so intensity and environmental sustainability have not been assessed for the same sites from EO. This review treats aspects of sustainability and intensification separately to provide a comprehensive overview of current research and the potential contribution of Earth Observation. The next two sections provide a more detailed overview of the methods used to assess both agricultural intensity and sustainability, highlighting the ways in which satellite data is used at present and the opportunities moving forwards.

3. Assessing the Intensity of Arable Production

3.1 Current Approaches

The types of indicators used to assess agricultural intensity differ between studies. Some focus on indicators which reflect the increase in land productivity caused by human intervention (Dietrich *et al.*, 2012) such as yield per ha (e.g. Singh *et al.*, 2002). Others focus on indicators which measure the change in inputs or other factors of management (Shriar, 2000) including total nitrogen (fertiliser) input (e.g. Temme and Verburg, 2011), number of pesticide applications (e.g. Herzog *et al.*, 2006), and inputs costs per ha (e.g. Teillard *et al.*, 2012). As such, the range of indicators used to assess agricultural intensity can be split into two general groups: agricultural input indicators (e.g. input cost per ha, crop acreage) and agricultural output indicators (e.g. production per area and time). Examples of indicators used in EO-based studies can be found in table 1. Some direct indicators of agricultural intensity, such as fertiliser and pesticide input cannot be measured using EO, but may be detectable indirectly from, for example, changes in yield. Typically either a single indicator is adopted to assess intensity (e.g. Biradar and Xiao, 2011; Mingwei *et al.*, 2008), or multiple indicators are aggregated to produce an intensity index (e.g. Kerr and Cihlar, 2003; Shriar, 2000). Aggregated indicators simplify complex situations into a single element, but this is done at the expense of interpretability and transparency. Whether a single indicator or an index is appropriate will vary depending on the purpose of the study (Herzog *et al.*, 2006).

Agricultural Intensity

- Crop yield (e.g. tonnes/ha)
- Multi-cropping: *Number of harvests within a single year (i.e. growing season)*
- Cropping intensity: *Number of cropping cycles per year or number of years a field is sown with crops and actually reaches harvest*
- Cropping frequency: *Number of years a pixel was cropped over an observation*
- Crop duration ratio: *Ratio of time period (during growing season) for which a pixel was cropped and the total length of the growing season*
- Fallow cycles: *Recurring periods of fallow cropland*
- Cropping area (e.g. acres, km²)

Further information in table S2 (supplementary material)

Table 1: Key EO-derived indicators for assessing agricultural intensity. *Examples of methods to derive these EO-based indicators can be found in tables S2 in the supplementary material.*

Data sources for agricultural intensity assessments include interviews, government statistics, field and farm surveys, aerial photographs and satellite data (see table S1 in the supplementary material for further examples of data sources and indicators used by various authors to investigate agricultural intensity). EO techniques are fairly common within this area, with a range of satellites appearing within the literature including MODIS, AVHRR, Landsat and Sentinel. EO-based methods used to investigate agricultural intensity vary depending on the indicator of interest; some examples of specific methods can be found in table S2 in the supplementary material.

EO has a range of advantages over other data sources for intensity assessments. The use of EO allows better cross-country comparisons to be made; country boundary restrictions do not apply in the same way as to government statistics data, thus improving data consistency (Herzog *et al.*, 2006). Additionally, low-cost methods using relatively simple technology can be developed, allowing application of EO-based methods in areas where costs of traditional data collection methods prohibit reliable intensity assessments (Ferencz *et al.*, 2004). Furthermore, EO-based cropping indicators perform well for broad-scale agricultural monitoring, suggesting they could complement (potentially) more accurate sample-based ground-data, by providing wall-to-wall observations of agricultural management (Estel *et al.*, 2016). Lack of spatially distributed information on key environmental and agronomic variables tends to limit application of crop simulation models for regional scale yield estimation (Moriondo *et al.*, 2007). EO data can alleviate this problem, by providing estimates of relevant variables over a range of spatial and temporal scales.

3.2 Potential for Expanding the Use of EO to Assess Agricultural Intensity

As previously noted, the use of EO within agricultural intensity studies is already fairly common, but it is still not typically routine and operational. In addition, with continued advancements in sensor technology, the launch of new satellites and the development of new methods, the full potential of EO has yet to be realised. Moving forward research should continue to focus on the creation of high resolution, global products which can be provided regularly (annually), consistently and in a timely manner (e.g. Claverie *et al.*, 2018; Egorov *et al.*, 2019; Roy *et al.*, 2010). In the past, the production of high resolution operational EO-derived products was hindered by a lack of suitable cloud-free imagery and the time and computing power required to process the vast number of images needed to provide global coverage. However now, with an increase in the number of moderate and high resolution satellites, improvements in gap-filling and sensor integration techniques, and the advent cloud computing systems that can facilitate more rapid processing, the potential for producing high resolution EO products for assessing agricultural intensity on a global scale has never been greater. Work is already underway to produce a variety of large-scale high-resolution (30m) products, for example, cropland extent maps on a country- (e.g. Teluguntla *et al.*, 2018) and continental-scale (e.g. Xiong *et al.*, 2017). International programmes such as GEOGLAM and commercial organisations such as OneSoil (OneSoil, 2018) are also working to map various agricultural parameters, including crop type and crop condition/development, on a global scale. The timely production of high resolution, global products, which provide an accurate representation of the diverse agrosystems around the world, is likely to require

either the development of generic, transferable models or an increase in the collection and provision of *in situ* data. In reality, the solution will probably involve some combination of the two.

The accessibility of appropriate field data for calibration and validation is a major constraint on the development of an operational EO-based system for assessing intensification. A step-change in monitoring capability could be provided by having EO as part of an integrated system that makes *in situ* data routinely available. Similarly, the creation and adoption of better data fusion methods may help with interpretation of the EO system, limiting the impact of confounding factors, and improving assessment potential. However, despite the various challenges which exist at present, EO data provides an important, practical and viable approach for regional and global monitoring of land surface dynamics, including variations in agricultural intensity (Yan *et al.*, 2014). The value of EO for agricultural intensity assessments lies, in part, in the spatially explicit nature of the data, consistency across political borders and the systematic acquisition setup (Kuemmerle *et al.*, 2013).

4. Assessing the Environmental Sustainability of Arable Systems

4.1 Current Approaches

Sustainability has three dimensions: economic, social and environmental sustainability (Allahyari *et al.*, 2016). EO has the potential to contribute valuable information on the environmental dimension however, to provide a comprehensive assessment of sustainability, EO must be used alongside datasets covering economic and social variables. Accepted socio-economic indicators are regularly (e.g. yearly) monitored by governments and international organisations (e.g. United Nations, World Health Organisation) on a regional, national and global scale. This provides a readily available, long-term dataset for economic and social sustainability, something that is not currently available for environmental sustainability. Consequently, in this review the focus is on the environmental dimension of sustainability.

Assessing environmental sustainability is a complex process fraught with challenges and pitfalls. Selection of appropriate indicators, weighting and aggregation methods for specific situations and requirements are essential to the successful assessment of sustainability. A number of frameworks have been developed to aid in the selection and aggregation of appropriate indicators, to provide a single score by which sustainability of agricultural systems can be assessed. These frameworks differ in their definition of sustainability, indicator selection approach, and aggregation and validation methods. Frameworks include

- SAFA: Sustainability Assessment of Food and Agriculture Systems (FAO, 2014, 2013)
- IDEA: Indicateur de Durabilité des Exploitations Agricoles (Zahm *et al.*, 2008)
- ISAP: Indicator of Sustainable Agricultural Practice (ISAP) (Rigby *et al.*, 2001)
- RISE: Response-Inducing Sustainability Evaluation (RISE) (Häni *et al.*, 2003, 2006)
- SAFE: Sustainability Assessment of Farming and the Environment (Van Cauwenbergh *et al.*, 2007)
- SSP: Sustainability Solution Space for Decision Making (Wiek and Binder, 2005)
- Sustainable Intensification Assessment Framework (Musumba *et al.*, 2017; Snapp *et al.*, 2018)

Issues surrounding the use of indicators and frameworks are explored in other publications (e.g. Binder *et al.*, 2010; Binder and Wiek, 2006; Gómez-Limón and Sanchez-Fernandez, 2010; Roy and Chan, 2012; Singh *et al.*, 2009; Smith *et al.*, 2017; Stein *et al.*, 2001) so no further discussion will be presented here.

Current agricultural sustainability studies generally rely on a mixture of primary and secondary data including: questionnaires, field data collection (e.g. soil sampling, spatial information) and government statistics (e.g.

Gómez-Limón and Sanchez-Fernandez, 2010; Rasul and Thapa, 2004; Rodrigues *et al.*, 2010; Zhen *et al.*, 2005). There are however a number of challenges specifically arising as a consequence of current data sources and collection methods which limit the ability to accurately and efficiently assess environmental sustainability:

- (1) **No baseline data** – Sustainability studies frequently lack baseline data, using data from a single point in time. This prevents analysis of temporal variability. Multi-temporal datasets would enable more comprehensive and therefore more reliable assessments.
- (2) **Uncertainty from sample data interpolation** – Reliance on sample data and interpolation or averaging techniques limit the potential to accurately assess spatial variability. Ideally all points would be sampled, providing (near-) continuous coverage, however, the cost and time required to complete such a task, clearly makes this impossible using traditional data collection techniques. Current assessments assume the sample data are representative of the wider study area, which may introduce error.
- (3) **Subjective data** – Frequent use of questionnaires, farm surveys and interviews arguably affects the objectivity of many studies. Apart from obvious issues over the truth of answers, subjectivity of qualitative collection methods limits the extent to which data can be integrated and compared. Use of objective, quantified data would likely improve assessment capabilities.
- (4) **Limited data** – Data availability, resolution and coverage (both spatial and temporal) are often limited by costly and time-consuming collection methods. Bias may also exist, with data less readily available in more inaccessible and poorer areas. Sustainability assessments are important in all areas if food demands are to be met and environmental quality maintained or improved; lack of necessary data due to traditional data collection methods hinders this.

EO has the potential to reduce some of these issues when assessing environmental sustainability.

4.2 Potential Applications of EO-based Methods for Assessing Environmental Sustainability

The potential applications for EO-based methods are described by splitting environmental sustainability into five key areas: (1) vegetation health; (2) soil quality; (3) water quality and availability; (4) biodiversity; and (5) ecosystem health (table 2). Environmental sustainability does encompass other aspects however the ability to assess elements such as air quality using EO, at scales relevant to agricultural systems, is restricted. This review therefore focuses on the areas considered to have most potential for assessment via EO-derived indicators for agriculture-based studies. Research efforts have not been evenly split between the areas covered in this section. However, to maintain a consistent structure and attempt to provide a balanced overview, each has approximately equal coverage in this section.

Environmental Sustainability	
Vegetation health	
<ul style="list-style-type: none"> • Crop condition • Biophysical traits <i>inc. biomass, fraction of absorbed photosynthetically active radiation (fAPAR), photosynthetic activity</i> • Structural traits <i>inc. crop/canopy height, leaf area index (LAI), biomass, canopy morphology</i> • Biochemical traits <i>inc. chlorophyll (Ch), water content, nitrogen (N) and phosphorous (P)</i> 	
<i>Further information in table S3 (supplementary material)</i>	
Soil Quality	
<ul style="list-style-type: none"> • Soil organic carbon (SOC) • Soil organic matter (SOM) • Soil moisture content • Soil salinity • Crop residue/conservation tillage density • Nitrogen status/availability 	
<i>Further information in table S4 (supplementary material)</i>	
Soil erosion/protection	
<ul style="list-style-type: none"> • Vegetation cover • Erosion feature detection • Erosion modelling <i>e.g. USLE</i> 	
<i>Further information in table S4 (supplementary material)</i>	
Water Quality	
<ul style="list-style-type: none"> • Water Quality Indices <i>derived from different spectral band combinations</i> • Physical water quality parameters <i>inc. total suspended solids (TSS), turbidity, suspended sediment concentration (SSC), chlorophyll concentration, temperature and water clarity</i> • Chemical water quality parameters <i>inc. concentration of total nitrogen, NO₃-N (nitrate as nitrogen) and total phosphorous</i> • Water quality proxy <i>e.g. health of vegetation alongside water bodies</i> 	
<i>Further information in table S5 (supplementary material)</i>	
Water Availability	
<ul style="list-style-type: none"> • Water body area and configuration • Water use efficiency and crop water stress • Water level and volume 	
<i>Further information in table S5 (supplementary material)</i>	
Biodiversity	
<ul style="list-style-type: none"> • Direct mapping of individuals and associations • Plant (and animal) species diversity • Habitat suitability <i>based on known habitat requirements of specific species</i> • Species Richness • Landscape structure <i>inc. composition, isolation and complexity</i> • Invasive Species 	
<i>Further information in table S6 (supplementary material)</i>	
Ecosystem Health	
<ul style="list-style-type: none"> • Vigour <ul style="list-style-type: none"> - Net Primary Productivity (NPP) & Gross Primary Productivity (GPP) - Fractional cover of green vegetation, non-photosynthetic vegetation (NPV) and bare soil - Biochemical properties <i>inc. nitrogen, phosphorous and chlorophyll</i> • Organisation <ul style="list-style-type: none"> - Species richness and biodiversity - Vegetation structural traits • Resilience • Ecosystem Services as a Proxy for Ecosystem Health 	
<i>Further information in table S7 (supplementary material)</i>	

Table 2: Potential EO-based indicators which could be used to assess environmental sustainability. *Examples of EO-based methods to derive these indicators can be found in tables S3 to S7 in the supplementary material.*

4.2.1 Vegetation Health

Environmental quality depends in part on the presence of healthy, diverse and abundant vegetation to provide ecosystem services including soil protection, carbon sequestration and flood prevention (Crossman *et al.*, 2013; Hein, 2014). Vegetation health, in turn, relies on a healthy environment to provide essential resources, including stable soil substrate and nutrients. This interdependence suggests that agricultural and non-agricultural vegetation health can be used as an indicator of environmental quality within the agricultural system. Various aspects of vegetation health have been assessed using EO over a range of spatial (e.g. field-/plot-scale to tens/hundreds km² and global scale) and temporal scales (e.g. single date assessments to decadal variation), in a

diverse range of environments (e.g. grasslands, shrublands, forests, rainforests, mountainous regions), including agricultural systems (e.g. corn farms, irrigated maize). Many of these studies were conducted using freely available satellite data including Landsat, MODIS and AVHRR, suggesting established methods exist, which can be readily applied to a range of environments. Examples of EO-based methods used to assess vegetation health can be found in table S3 in the supplementary material.

Empirical models are commonly used to assess a variety of vegetation health-related properties. The frequency and timing of image acquisition affects the strength of relationships between specific vegetation indices (VIs) (e.g. NDVI) and related variables (e.g. net primary productivity) (Tebbs *et al.*, 2017). In some situations it may therefore be preferable to use coarser spatial resolution satellites which provide daily coverage (e.g. MODIS), instead of finer spatial resolution satellites with less frequent data acquisition (e.g. Landsat) (Jackson *et al.*, 2004). The successful application of Radiative Transfer Models (RTMs) demonstrates the potential to develop (simple) algorithms to predict various plant traits from satellite data, spanning a range of vegetation types (Myneni *et al.*, 1997; Trombetti *et al.*, 2008).

4.2.2 Soil Quality

Deterioration of soil quality through intensive use of agricultural land has far-reaching impacts affecting plant productivity, water and air quality (Doran and Zeiss, 2000). Soil quality is therefore an important indicator of sustainability and has been studied over a range of scales, from 10s to 1000s km², and across diverse landscapes including cultivated, semi-natural and natural vegetation areas using EO data. Previous studies clearly demonstrate the great potential for soil quality assessment in agricultural areas. However, disparity between achievable spatial resolutions with current freely available satellite data (e.g. Landsat) and typical agricultural field sizes limits the ability to conduct field-scale assessments in some parts of the world. Examples of methods used to assess soil quality and erosion/protection using EO data can be found in table S4 in the supplementary material.

Soil reflectance is a function of the soil's physical properties such as soil moisture and soil organic carbon, but also tillage practices, crop residue and row orientation. Multispectral imagery is therefore best suited for application to farms with uniformly tilled fields and constant soil moisture conditions at the time of image acquisition; such conditions increase the dominance of the property of interest in the spectral response (Barnes and Baker, 2000). Microwave sensors allow direct soil moisture estimates by exploiting the relationship between moisture content and the dielectric constant of the soil (Wagner *et al.*, 2007). Polarisation and study site conditions influence detection of soil parameter variation. For example, at low moisture conditions, vertical polarisation offers higher sensitivity to salinity; at high moisture levels horizontal polarisation exhibits slightly higher sensitivity (Lasne *et al.*, 2008; Shoshany *et al.*, 2013).

4.2.3 Water Quality and Availability

AI can negatively affect both quality and availability of water resources, so sustainability assessments should consider both. EO-based water quality and availability assessments have focused predominantly on large water bodies such as lakes, reservoirs and coastal environments, for non-agriculture-related investigations. Although the methodology may theoretically be transferable to agricultural environments, the applicability of EO methods to sustainability assessments depends partly on the scale of the study. Water bodies of interest for field- or farm-based assessments are likely to be much smaller than can easily be detected by current satellites due to limiting spatial resolutions; as sensors continue to develop the potential to adopt such methods will increase. At present, EO-based water-related assessments are likely to be most suitable for catchment- or regional-scale investigations, where the impact of multiple farms on larger water bodies is considered using coarser resolution data. Some of the common EO-based methods used to assess water quality and availability are outlined in table S5 in the supplementary material.

Water quality assessments commonly employ empirical models to estimate physical and chemical water quality parameters. Transference of empirical methods to areas and circumstances for which they were not formulated adds uncertainty and may not be appropriate. For example, current generation satellite sensors have limited ability to map chlorophyll content in mesotrophic and oligotrophic water, despite successful application for eutrophic and hyper-eutrophic water (Gons *et al.*, 2008). However, understanding the limitations of the empirical relationships means they can be applied appropriately. For example a set of robust relationships established between suspended sediment concentration and MODIS spectral reflectance data can be applied, but only for rivers larger than 500m (Martinez *et al.*, 2008).

The operational feasibility of applying a single water body detection algorithm over diverse environmental and climatic conditions has been demonstrated, producing results with a high-degree of accuracy (Mueller *et al.*, 2016). The spatial resolution of satellite altimetry is about 1.7 to 3km (for calm waters), so water level assessments have typically targeted large lakes or rivers, however, satellite altimetry-based techniques have been successfully applied to medium-sized water bodies (200-800m wide), with some potential for application to small-sized water bodies (40-200m wide) (Sulistioadi *et al.*, 2015).

4.2.4 Biodiversity

AI has been linked with biodiversity losses on local, regional and global scales. Biodiversity helps maintain ecosystem health and provision of ecosystem services, so its loss can have serious implications for human well-being (Pettorelli *et al.*, 2014; Turner *et al.*, 2015). Preservation or ideally enhancement of biodiversity is essential for sustainable agriculture, hence the spatial and temporal variability of biodiversity is an important indicator of sustainability. The use of EO to monitor biodiversity has been reviewed by a number of authors, who have outlined key methods, challenges and opportunities (e.g. Gillespie *et al.*, 2008; Kerr and Ostrovsky, 2003; Kuenzer *et al.*, 2014; Mairota *et al.*, 2015; Nagendra, 2001; Rocchini *et al.*, 2010; Turner *et al.*, 2003; Wang *et al.*, 2010). Consequently, in this review we provide only a brief overview covering examples of indicators and related methods (table 2). Examples of some EO-based methods used to assess biodiversity can be found in table S6 in the supplementary material.

Biodiversity can be assessed using both direct and indirect techniques. Individual plants or associations of single species may be directly mapped using (very) high spatial resolution satellite images. Where direct mapping of biodiversity indicators is not feasible, indirect approaches that rely on environmental parameters as proxies may be adopted (Turner *et al.*, 2003). Both species abundance and richness are considered to be fundamentally affected by landscape heterogeneity; more heterogeneous landscapes can host more diverse species than homogeneous landscapes due to greater niche availability (Feng *et al.*, 2010; Honnay *et al.*, 2003). EO-derived measures of landscape structure such as composition, complexity and isolation can therefore be used to predict species distribution and diversity. Habitat fragmentation and removal of connectivity elements (e.g. hedgerows) negatively affect species richness and distribution; this relationships allows species richness predictions based on relatively simple landscape metrics (Griffiths and Lee, 2000; Honnay *et al.*, 2003). Lack of consideration of environmental factors such as temperature and disturbance, which affect biodiversity, may reduce accuracy of LC- and landscape metric-based predictions (Griffiths and Lee, 2000; Li *et al.*, 2014).

Satellite-derived habitat maps can be used in conjunction with known habitat requirements to model potential distribution and abundance of individual species (e.g. Nagendra, 2001; Weiers *et al.*, 2004); habitat suitability parameters such as the existence of suitable water bodies can be derived to assist this modelling (e.g. Weiers *et al.*, 2004). Habitat-based species distribution modelling requires LC data of sufficient spatial and thematic resolution to ensure all habitats that the target species could potentially occupy are identified; *in situ* or ancillary data is almost certainly necessary to meet requirements for predicting actual distributions of many species (Kerr and Ostrovsky, 2003). Where species do not occupy all suitable habitats, only potential rather than actual distribution

can be predicted (Davis *et al.*, 2007; Kerr and Ostrovsky, 2003). A limitation of using habitat maps to predict species distribution and diversity is the fact that this provides no insight into within-habitat variation (Nagendra, 2001).

4.2.5 Ecosystem Health

Ecosystem (ES) health affects the ability of ecosystems to provide essential services. A sustainable ecosystem is one which has the ability to maintain its function (or vigour) and structure (or organisation) over time and is resilient even with the application of external stress (Costanza and Mageau, 1999). The potential of EO has been acknowledged and reviewed previously by other authors (e.g. Andrew *et al.*, 2014; Feng *et al.*, 2010; Li *et al.*, 2014). Assessments generally focus on three aspects of ES health: *vigour*, a measure of a system's activity, metabolism and primary productivity; *organisation*, the number and diversity of interactions between a system's components; and *resilience*, a system's ability to maintain its pattern and structure in the presence of stress (Costanza and Mageau, 1999; Li *et al.*, 2014; Li and Guo, 2012). Common indicators of ES health are GPP or NPP (for vigour), species richness and diversity (for organisation) and the ratio of an ES health indicator pre- and post-disturbance (for resilience) (Costanza and Mageau, 1999; Li *et al.*, 2014; Li and Guo, 2012) (table 2). Examples of some possible EO-based methods for assessment of ecosystem health can be found in table S7 in the supplementary material.

Ecosystem health can also be assessed using the supply of ecosystem services (ESS) as a proxy. ESS represent the goods and services supplied by organisms and their activities, controlled by the abiotic characteristics of the system and the anthropogenic impacts it experiences (Andrew *et al.*, 2014). Agricultural systems are both a source and beneficiary of ESS (Balbi *et al.*, 2015; Power, 2010). RS-based ESS mapping has been reviewed by various authors (e.g. Andrew *et al.*, 2014; Crossman *et al.*, 2013). ESS supply has been mapped at various scales (e.g. sub-national to global) using both direct and indirect techniques. Whether direct or indirect techniques should be used depends on the information needs and characteristics of the ESS of interest (Andrew *et al.*, 2014). EO-based methods can improve ESS supply estimates through their ability to depict subtle spatial variations in the plant functional traits and soil properties known to influence the supply of many services (Andrew *et al.*, 2014).

It is worth noting that apart from microwave imagery, EO has limited capability for penetrating the vegetation canopy to extract useful information about any lower strata such as herbs or shrubs when the overstory is dense (Nagendra, 2001). Similarly, EO is incapable of providing direct information on below ground components of the ecosystems; these must be inferred from above ground data (Feng *et al.*, 2010). EO-based assessments may therefore only provide a partial picture of diversity and soil-based ESS, for example, restricting the potential to comprehensively assess ESS provision. This restriction of EO data highlights the importance of creating an integrated framework of EO and *in situ* data collection, not only to provide data for training and validation of EO products, but also to provide information on the areas that cannot be adequately assessed using EO.

5. Discussion

When using satellite data there are a number of considerations to take into account, including data gaps and image frequency. The availability, or lack thereof, of cloud-free images at suitable timescales is a critical challenge for many EO-based applications. Research suggests revisit frequencies ranging from <1 day to exactly 8 days (depending on location and time of year) would be necessary to achieve a view at least 70% clear within 8 days (Whitcraft *et al.*, 2015). Previously, such revisit times were only achievable using coarse resolution data, which restricted the ability to perform fine-scale assessments of agricultural intensity. However, the number of medium-high resolution satellites has increased in recent years, for example with the launch of those within the Copernicus Programme. Combining freely available data from these various systems increases the number of images available for each point on earth. For example, by combining Landsat 8 with Sentinel-2A/B data, one image should be available every 3 days on average (Li and Roy, 2017), increasing the likelihood of obtaining sufficient cloud-free images. The creation of a Harmonised Landsat and Sentinel-2 reflectance data set (Claverie *et al.*, 2018) alongside the free availability of

Sentinel-1 data (Torres *et al.*, 2012), which is not affected by cloud cover, the launch of satellite fleets (e.g. Planet Labs) and the increased use of UAVs are all helping to mitigate the impact of cloud-cover on EO-based applications.

Various techniques also exist to deal with gaps and noise in the data. These techniques include pixel unmixing (Zhang *et al.*, 2017), data fusion (Gevaert and García-Haro, 2015; Senf *et al.*, 2015; Wang and Atkinson, 2018), best-pixel selection (Griffiths *et al.*, 2013; Hermosilla *et al.*, 2018), data interpolation (Inglada *et al.*, 2017; Vuolo *et al.*, 2017), climatology fitting (Verger *et al.*, 2013), temporal smoothing (Kandasamy *et al.*, 2013; Shao *et al.*, 2016; Tan *et al.*, 2011) and temporal aggregation (Loveland *et al.*, 2000). The introduction of computing platforms such as Google Earth Engine (Gorelick *et al.*, 2017) allows these complex algorithms to be applied to large volumes of data, by providing access to greater computing power and satellite datasets on a global-scale (Carrasco *et al.*, 2019). These platforms greatly reduce the time and cost associated with image processing, which increases the viability of such gap-filling techniques and facilitates development of new approaches for producing necessary cloud-free datasets.

Another key issue is the development of universally applicable EO-based monitoring techniques. Methods, such as crop yield models, that rely on largely location- and sensor-specific empirical relationships to retrieve indicators are common within many EO-based applications. However, lack of historic and large-scale *in situ* data for all areas limits the potential for calibration and validation of such methods (Doraiswamy *et al.*, 2005; Estel *et al.*, 2016). Suitable validation and calibration data may exist, but availability to researchers may be restricted due to commercial confidentiality, among other factors. The frequent reliance on empirical relationships, and lack of *in situ* data for all locations, limits the transportability of these methods to different sensors and study areas (Andrew *et al.*, 2014; Li *et al.*, 2014); development of more generalised models is required. Less empirical models still often have parameters that need calibrating, to enable them to be applied appropriately to new geographical areas, new crops, or new animal species.

The creation of generic, transferable and widely applicable models must consider a variety of factors including the scale and resolution over which specific relationships apply (Foody, 2004; Gillespie *et al.*, 2008) (Gustafson, 1998) and the variation in the relative dominance of different variables within space and time (e.g. Prasad *et al.*, 2006). Additionally, the impact that landscape characteristics can have on the accuracy with which indicators can be retrieved needs to be considered. For example, the size, shape and orientation of objects, such as narrow rivers (Sulistioadi *et al.*, 2015) and small habitat patches (Luoto *et al.*, 2002), can limit their detectability. Underlying environmental conditions can also affect the accuracy with which variables such as crop residue (Pacheco and McNairn, 2010) and soil moisture (Lakhankar *et al.*, 2009) can be retrieved. As such, comparison over diverse agricultural environments must be done with care, and with an understanding of the underlying differences in landscape characteristics.

To create models that can be widely applied, either empirical models are required that are integrated with ground collection efforts (e.g. Boryan *et al.*, 2011), or model-based methods that need no calibration to be applied to new areas must be developed. In practice, the solution is likely to involve a mixture of these two options.

5.1 Opportunities for an EO-based SI Assessment System

Despite the potential importance of SI for securing future food supplies, methods for assessing the success of SI attempts are currently lacking. Gaps in these assessments arise, in part, due to the reliance on data that typically lacks the temporal and spatial coverage and resolution necessary to make a reliable assessment of SI. EO provides a range of opportunities for the development of an operational SI monitoring system that can be applied from field-scale to global-scale, at various temporal resolutions. The global coverage and consistency (e.g. image resolution, data quality and processing standards) afforded by EO data facilitates multi-scale analysis and comparisons between countries with diverse farming practices and field data availability. Additionally, the ability to derive different indicators from the same EO data sources ensures coherence between measures of sustainability

and intensification; this facilitates easier, more reliable integration of indicators into an assessment framework. The availability of long-term data (30-40 years from Landsat data archive) and repeat measurements allows the establishment of a baseline against which long-term and short-term changes can be assessed. Combined with the increasing availability of free satellite data, these factors allow a more flexible, adaptable and cost-effective approach to SI assessment to be developed. Furthermore, the ease, speed and efficiency with which assessments can be conducted is increased by the digital nature of EO data which allows, for example, simpler data input and processing. Recent advances in computer processing power/capability, such as cloud computing, further improve the situation, enhancing our ability to process and analyse large datasets.

As this review demonstrates, the basic EO-based assessment techniques for indicators of environmental sustainability and agricultural intensity are already established. This means that future work can focus on amalgamating existing work and creating a framework to integrate relevant indicators. An important step in the development of a comprehensive EO-based assessment framework is the creation of a set “Essential Sustainable Intensification Variables” (ESIVs) to form the basis for a global monitoring program. Having a set of “essential variables” helps to prioritise efforts by outlining a minimum set of essential measurements (Pereira *et al.*, 2013) required to capture major dimensions of agricultural and environmental change, allowing the sustainability of intensification to be assessed. The development of this list should build on the selection frameworks created for the Essential Climate Variables (ECVs) and Essential Biodiversity Variables (EBVs) of the Global Framework for Climate Services (GCFS) and Group on Earth Observations Biodiversity Observation Network (GEO BON) respectively. In brief, variable selection will require an open, inter-disciplinary process, involving the engagement of scientific, policy and other communities (Pereira *et al.*, 2013). This will ensure it builds on existing activities such as the ECVs, EBVs and GEOGLAM. Variables should be identified based on key criteria such as relevance, feasibility, scalability, temporal sensitivity and cost effectiveness (Bojinski *et al.*, 2014; Pereira *et al.*, 2013). As Pettorelli *et al.* (2016) highlight for the ECVs and EBVs, the identification of suitable ESIVs will be an evolving process. The list of indicators will need to be periodically updated as technology advances, and as sensor availability and observation priorities change. Table 3 provides an example list of possible “essential” EO-based indicators. This list is based on indicators already incorporated in the EBVs and ECVs, and previous SI frameworks including the Sustainable Intensification Assessment Framework (Musumba *et al.*, 2017).

ESIV examples	Relevance	Existing Operational Products	Ideal Product Coverage
Agricultural intensity			
Crop type	Essential product required to be able to accurately monitor/derive crop yield and area, but not an indicator in its own right.	<ul style="list-style-type: none"> • OneSoil crop-type map 2016, 2017, 2018 • Country-level products e.g. CEH Land Cover plus: Crops 2015 (partial coverage), 2016, 2017, 2018 (GB) 	<ul style="list-style-type: none"> • Annually (possibly more often for multi-cropping systems)
Crop yield	Needed to quantify agricultural intensity	Currently no operational products exist	<ul style="list-style-type: none"> • Annually
Crop yield gap	Needed to help identify areas that could be farmed more intensively	Currently no operational products	<ul style="list-style-type: none"> • Annually
Cropping area	Needed to quantify agricultural extent	<ul style="list-style-type: none"> • OneSoil crop-type map 2016, 2017, 2018 Crops are a subset of land cover maps: <ul style="list-style-type: none"> • Country-specific Land Cover maps e.g. CEH GB Land Cover Map 2015 30m • Pan-European: CORINE Land Cover minimum mapping unit 25ha • Global land cover map 30m (Chen <i>et al.</i>, 2015) 	<ul style="list-style-type: none"> • Annually

Environmental sustainability			
NPP/GPP*	Provides a measure of the health/degradation of the ecosystem; underpins all production-based ecosystem services.	<ul style="list-style-type: none"> • NASA MODIS yearly 500m/1km 2000 to present • Copernicus GDMP 10-day 1km 1999 to present & 10-day 300m 2014 to present 	<ul style="list-style-type: none"> • Patches of natural/semi-natural habitat within the farmed area • Annually
Soil moisture ×	Indicator of soil quality. Increased fertiliser use can increase water consumption and deplete soil moisture. A decline in soil moisture may be an indication of unsustainable intensification.	<ul style="list-style-type: none"> • SMOS daily/3-day/10-day/monthly 15/25km 2009 to present • Copernicus METOP/ASCAT daily 0.01° 2007 to present • Copernicus Sentinel-1 daily 1km 2015-2017 (Europe only) 	<ul style="list-style-type: none"> • Farmed area • Annually
Soil erosion	Removal of interstitial features (e.g. hedgerows) and increased runoff (due to soil compaction) increases soil erosion. An increase in soil erosion may be an indication of unsustainable intensification.	Currently no operational products exist	<ul style="list-style-type: none"> • Farmed area plus surrounding area • Annually
Soil organic carbon ×	Indicator of soil quality. Increased irrigation and soil erosion lead to a decline in organic matter content. A decrease in soil organic carbon may be an indication of unsustainable intensification.	Currently no operational products exist	<ul style="list-style-type: none"> • Farmed area • Annually
Water clarity/turbidity	Increased agrochemical inputs and increased soil erosion reduce water quality. Lower clarity/higher turbidity may be an indication of unsustainable intensification.	<ul style="list-style-type: none"> • Copernicus ENVISAT/MERIS 10-day 300m/1km 2002 to 2012 • Copernicus Sentinel-3/OLCI 10-day 300m/1km 2017 to present 	<ul style="list-style-type: none"> • Nearby water bodies • Annually
Landscape structure*	Removal of interstitial features e.g. hedges and increased field sizes cause simplification of habitat structure and loss of ecosystem connectivity. Knock-on effect on species populations and diversity (e.g. birds).	<ul style="list-style-type: none"> • Country-specific Land Cover maps e.g. CEH GB Land Cover Map 2015 30m • Pan-European: CORINE Land Cover Minimum mapping unit 25ha • Global land cover map 30m (<i>Chen et al., 2015</i>) 	<ul style="list-style-type: none"> • Surrounding area • Annually
Species richness/diversity*	Increase in agrochemical inputs, irrigation and landscape structure simplification alter species composition. A decrease in species richness and diversity may be an indication of unsustainable intensification.	<ul style="list-style-type: none"> • European Atlas of Forest Tree Species 1km 	<ul style="list-style-type: none"> • Surrounding area • Annually • Flora and fauna species

*Table 3: Examples of possible Essential Sustainable Intensification Indicators (ESIV). *indicates variables which are already included within the list of EBVs. × indicates variables which are already included within the list of ECVs. NB: the OneSoil data is not currently available for download, but it shows the potential for the creation of an operational global crop-type map.*

Once a comprehensive list of ESIVs has been generated, careful consideration must be given to the selection of appropriate methods to assess each variable. This could be achieved by adopting an open process of algorithm inter-comparison and selection similar to that used by ESA in their Climate Change Initiative (Hollmann *et al.*, 2013), 2013) and the Sen2-Agri project (Bontemps *et al.*, 2015). To allow reliable comparison of different algorithms and to ensure their relevance at local scales and widely varying agricultural systems at global scales, an open test dataset similar to that used in the Sen2-Agri project should be developed. This dataset was created through acquisition of satellite and *in situ* data from the same season over sites representative of global agricultural system diversity (Bontemps *et al.*, 2015). Application of the potential algorithms to this data set ensured an objective and transparent algorithm selection method, which should be mirrored in the creation of the ESIV data products.

An important part of building the ESIV products will be a comprehensive assessment of the associated uncertainties and a clear communication of these uncertainties to the end-user. Kissling *et al.* (2018) set out a workflow of 11 steps used to operationalise the process of building EBV data products, including the quantification and communication of uncertainties in terms of data, model algorithms and parameters. Consideration must be

given to uncertainties associated with the underlying raw data, from both satellite and *in situ* sources, and from processing methods (e.g. gap-filling techniques) and models applied to this data. Kissling *et al.* (2018) highlight the need to develop high-throughput processing tools for quantifying uncertainties; the same will be true for the ESIVs.

Another key element will be the development of a framework through which the ESIVs can be utilised to provide an assessment of the sustainability of agricultural intensification. Previous assessment frameworks have utilised indicators in a number of ways: (i) individually, expressed in units, (ii) as part of a set, or (iii) in a composite index, whereby scores of individual indicators are combined into a single, dimensionless number, or sustainability score (Dantsis *et al.*, 2010; Farrell and Hart, 1998; Mitchell *et al.*, 1995; Van Passel and Meul, 2012). An example of a widely used composite indicator for sustainability assessment is the Ecological Footprint which combines various indicator footprints including carbon, forest, crop land, & built up land footprint to provide a measurement of human demand for land and water areas (Toderiou, 2010 in Čuček *et al.*, 2012; Galli *et al.*, 2012). Some studies choose not to aggregate their indicators, adopting instead the use of sustainability polygons, webs and radars, which removes the need for aggregation across different scales by displaying scores for different index components simultaneously (Rigby *et al.*, 2001). A decision about the best way to utilise the ESIVs within the proposed EO-based assessment framework will require consultation with researchers and intended users.

The success of the proposed EO-based assessment framework will rely heavily on the development of an integrated system of routine collection and provision of *in situ* data. *In situ* data is required to perform a number of roles including calibration and validation of EO-derived products and assessment of elements of the environment that cannot efficiently/effectively be monitored using EO data, for example below-ground properties and processes. A comprehensive assessment of SI will also require economic and social data that cannot be provided by EO.

An EO-based assessment framework could be implemented at different scales and at different levels of detail. For example at country-level, assessments are more likely to have access to environmental and farming data sets that would enable more detailed assessments of environmental sustainability. National-scale assessments are also more likely to have access to additional economic and social data that would enable more comprehensive SI assessments to be conducted. However, globally less detailed *in situ* data is likely to be available, although programs such as GEOSS and GEOGLAM have shown that good quality reference data sets can be collected for some areas to help the development of more global solutions. The development of detailed nationally-based methods, and less detailed globally-based methods, is likely to occur in different ways. A globally-based system would develop most sensibly through integration with existing global initiatives such as GEOSS, which is already generating products capable of agricultural and environmental monitoring. Whereas national-scale solutions are likely to develop from existing country-level environmental and agricultural monitoring schemes. However, for both the global and national-scale, the more integrated the EO and other strands of environmental monitoring are the better system will be.

6. Conclusion

One element of meeting the future food demands of a growing population, with shifting food consumption patterns, will be the intensification of agricultural production. To ensure long term environmental degradation is avoided, any increases in food production must be undertaken in a sustainable manner. The lack of any prescribed methods of sustainable intensification mean that to successfully achieve this goal a comprehensive method of assessing the sustainability of intensification endeavours must be developed. Various frameworks exist at present; however, these commonly rely on traditional data sources that do not provide adequate coverage, resolution, or frequency of data to generate reliable results for all agricultural systems. The potential for an EO-based assessment system is clear, with an extensive body of research into EO methods for monitoring earth surface properties and their spatial and temporal variation.

The element that is currently missing is a system for combining these indicators to provide a comprehensive assessment of the sustainability of agricultural intensification. Such a system could build on the approaches used to develop the EBVs and ECVs and global agricultural monitoring schemes such as GEOGLAM. Determining the optimum format for this system will require a multi-disciplinary, multi-organisation working group involving farmers, researchers, government bodies and other stakeholders.

Irrespective of the exact nature of the final system, EO offers the opportunity to obtain more spatially and temporally representative data, over scales and resolutions unobtainable with conventional data collection methods. An EO-based system, however, does not exclude the need for *in situ* data, rather it will supplement current systems facilitating more efficient and consistent multi-scale assessments over a range of temporal resolutions at a lower cost. Integration of EO and *in situ* data on national and global scales, will provide a step change in our ability to provide regular, consistent and timely assessments. This is essential if we are to meet future production demands without causing significant, irreparable damage to the environment.

Acknowledgements

MH was funded by Lancaster University through a Lancaster Environment Centre PhD Studentship as part of the Graduate School for the Environment. CR was supported by research programme NE/N018125/1 LTS-M ASSIST - Achieving Sustainable Agricultural Systems, funded by NERC and BBSRC.

Declarations of interest: none

References

- Allahyari, M.S., Masouleh, Z.D., Koundinya, V., 2016. Implementing Minkowski fuzzy screening, entropy, and aggregation methods for selecting agricultural sustainability indicators. *Agroecol. Sustain. Food Syst.* 40, 277–294. doi:10.1080/21683565.2015.1133467
- Andrew, M.E., Wulder, M.A., Nelson, T.A., 2014. Potential contributions of remote sensing to ecosystem service assessments. *Prog. Phys. Geogr.* 38, 1–26. doi:10.1177/0309133314528942
- Balbi, S., del Prado, A., Gallejones, P., Geevan, C.P., Pardo, G., Pérez-Miñana, E., Manrique, R., Hernandez-Santiago, C., Villa, F., 2015. Modeling trade-offs among ecosystem services in agricultural production systems. *Environ. Model. Softw.* 72, 314–326. doi:10.1016/j.envsoft.2014.12.017
- Barnes, A.P., Thomson, S.G., 2014. Measuring progress towards sustainable intensification: How far can secondary data go? *Ecol. Indic.* 36, 213–220. doi:10.1016/j.ecolind.2013.07.001
- Barnes, E.M., Baker, M.G., 2000. Multispectral data for mapping soil texture: possibilities and limitations. *Appl. Eng. Agric.* 16, 731–741. doi:10.13031/2013.5370
- Benton, T.G., Vickery, J.A., Wilson, J.D., 2003. Farmland biodiversity: Is habitat heterogeneity the key? *Trends Ecol. Evol.* 18, 182–188. doi:10.1016/S0169-5347(03)00011-9
- Binder, C.R., Feola, G., Steinberger, J.K., 2010. Considering the normative, systemic and procedural dimensions in indicator-based sustainability assessments in agriculture. *Environ. Impact Assess. Rev.* 30, 71–81. doi:10.1016/j.eiar.2009.06.002
- Binder, C.R., Wiek, A., 2006. The role of transdisciplinary processes in sustainability assessment of agricultural systems, in: Häni, F.J., Pintér, L., Herren, H.R. (Eds.), *Sustainable Agriculture: From Principles to Common Practice. Proceedings and Outputs of the First Symposium of the International Forum on Assessing Sustainability in Agriculture (INFASA)*, March 16, 2006, Bern, Switzerland. International Institute for Sustainable Development, pp. 33–48.
- Bingfang, W., 2006. Introduction of China Crop Watch system with remote sensing, in: *ISPRS Archives*. XXXVI-8/W48:

- Remote Sensing Support to Crop Yield Forecast and Area Estimates. Stresa, Italy, pp. 15–18.
- Biradar, C.M., Xiao, X., 2011. Quantifying the area and spatial distribution of double-and triple-cropping croplands in India with multi-temporal MODIS imagery in 2005. *Int. J. Remote Sens.* 32, 367–386. doi:10.1080/01431160903464179
- Bojinski, S., Verstraete, M., Peterson, T.C., Richter, C., Simmons, A., Zemp, M., 2014. The Concept of Essential Climate Variables in Support of Climate Research, Applications, and Policy. *Bull. Am. Meteorol. Soc.* 95, 1431–1443. doi:10.1175/bams-d-13-00047.1
- Bommarco, R., Kleijn, D., Potts, S.G., 2013. Ecological intensification: Harnessing ecosystem services for food security. *Trends Ecol. Evol.* 28, 230–238. doi:10.1016/j.tree.2012.10.012
- Bontemps, S., Arias, M., Cara, C., Dedieu, G., Guzzonato, E., Hagolle, O., Inglada, J., Matton, N., Morin, D., Popescu, R., Rabaute, T., Savinaud, M., Sepulcre, G., Valero, S., Ahmad, I., Bégué, A., Wu, B., de Abelleira, D., Diarra, A., Dupuy, S., French, A., Akhtar, I. ul H., Kussul, N., Lebourgeois, V., Page, M. Le, Newby, T., Savin, I., Verón, S.R., Koetz, B., Defourny, P., 2015. Building a data set over 12 globally distributed sites to support the development of agriculture monitoring applications with Sentinel-2. *Remote Sens.* 7, 16062–16090. doi:10.3390/rs71215815
- Boryan, C., Yang, Z., Mueller, R., Craig, M., 2011. Monitoring US agriculture: The US department of agriculture, national agricultural statistics service, cropland data layer program. *Geocarto Int.* 26, 341–358. doi:10.1080/10106049.2011.562309
- Buckwell, A., Nordang Uhre, A., Williams, A., Poláková, J., H Blum, W.E., Schiefer, J., Lair, G.J., Heissenhuber, A., Schießl, P., Krämer, C., Haber, W., 2014. The Sustainable Intensification of European Agriculture. A Review Sponsored by the RISE Foundation.
- Campbell, B.M., Thornton, P., Zougmore, R., van Asten, P., Lipper, L., 2014. Sustainable intensification: What is its role in climate smart agriculture? *Curr. Opin. Environ. Sustain.* 8, 39–43. doi:10.1016/j.cosust.2014.07.002
- Carrasco, L., O'Neil, A.W., Daniel Morton, R., Rowland, C.S., 2019. Evaluating combinations of temporally aggregated Sentinel-1, Sentinel-2 and Landsat 8 for land cover mapping with Google Earth Engine. *Remote Sens.* 11, 288. doi:10.3390/rs11030288
- Caviglia, O.P., Andrade, F.H., 2010. Sustainable Intensification of Agriculture in the Argentinean Pampas: Capture and Use Efficiency of Environmental Resources. *Am. J. Plant Sci. Biotechnol.* 3, 1–8.
- Claverie, M., Ju, J., Masek, J.G., Dungan, J.L., Vermote, E.F., Roger, J.C., Skakun, S. V., Justice, C., 2018. The Harmonized Landsat and Sentinel-2 surface reflectance data set. *Remote Sens. Environ.* 219, 145–161. doi:10.1016/j.rse.2018.09.002
- Costanza, R., Mageau, M., 1999. What is a healthy ecosystem? *Aquat. Ecol.* 33, 105–115. doi:10.1023/A:1009930313242
- Crossman, N.D., Burkhard, B., Nedkov, S., Willemen, L., Petz, K., Palomo, I., Drakou, E.G., Martín-Lopez, B., McPhearson, T., Boyanova, K., Alkemade, R., Egoh, B., Dunbar, M.B., Maes, J., 2013. A blueprint for mapping and modelling ecosystem services. *Ecosyst. Serv.* 4, 4–14. doi:10.1016/j.ecoser.2013.02.001
- Crowder, D.W., Jabbour, R., 2014. Relationships between biodiversity and biological control in agroecosystems: Current status and future challenges. *Biol. Control* 75, 8–17. doi:10.1016/j.biocontrol.2013.10.010
- Čuček, L., Klemeš, J.J., Kravanja, Z., 2012. A review of footprint analysis tools for monitoring impacts on sustainability. *J. Clean. Prod.* 34, 9–20. doi:10.1016/j.jclepro.2012.02.036
- Dantsis, T., Douma, C., Giourga, C., Loumou, A., Polychronaki, E.A., 2010. A methodological approach to assess and compare the sustainability level of agricultural plant production systems. *Ecol. Indic.* 10, 256–263. doi:10.1016/j.ecolind.2009.05.007
- Davis, F.W., Seo, C., Zielinski, W.J., 2007. Regional variation in home-range-scale habitat models for fisher (*Martes*

- pennanti) in California. *Ecol. Appl.* 17, 2195–2213. doi:10.1890/06-1484.1
- Dietrich, J.P., Schmitz, C., Müller, C., Fader, M., Lotze-Campen, H., Popp, A., 2012. Measuring agricultural land-use intensity - A global analysis using a model-assisted approach. *Ecol. Modell.* 232, 109–118. doi:10.1016/j.ecolmodel.2012.03.002
- Dillon, E.J., Hennessy, T., Buckley, C., Donnellan, T., Hanrahan, K., Moran, B., Ryan, M., 2016. Measuring progress in agricultural sustainability to support policy-making. *Int. J. Agric. Sustain.* 14, 31–44. doi:10.1080/14735903.2015.1012413
- Dillon, E.J., Hennessy, T., Hynes, S., 2010. Assessing the sustainability of Irish agriculture. *Int. J. Agric. Sustain.* 8, 131–147. doi:10.3763/ijas.2009.0044
- Donald, P.F., Green, R.E., Heath, M.F., 2001. Agricultural intensification and the collapse of Europe's farmland bird populations. *Proc. R. Soc. B* 268, 25–29. doi:10.1098/rspb.2000.1325
- Doraiswamy, P.C., Sinclair, T.R., Hollinger, S., Akhmedov, B., Stern, A., Prueger, J., 2005. Application of MODIS derived parameters for regional crop yield assessment. *Remote Sens. Environ.* 97, 192–202. doi:10.1016/j.rse.2005.03.015
- Doran, J.W., Zeiss, M.R., 2000. Soil health and sustainability: managing the biotic component of soil quality. *Appl. Soil Ecol.* 15, 3–11. doi:10.1016/S0929-1393(00)00067-6
- Egorov, A. V., Roy, D.P., Zhang, H.K., Li, Z., Yan, L., Huang, H., 2019. Landsat 4, 5 and 7 (1982 to 2017) Analysis Ready Data (ARD) observation coverage over the conterminous United States and implications for terrestrial monitoring. *Remote Sens.* 11, 447. doi:10.3390/rs11040447
- Elliott, J., Firbank, L.G., Drake, B., Cao, Y., Gooday, R., 2013. Exploring the Concept of Sustainable Intensification. ADAS/Firbank, LUPG Commissioned Report.
- Estel, S., Kuemmerle, T., Levers, C., Baumann, M., Hostert, P., 2016. Mapping cropland-use intensity across Europe using MODIS NDVI time series. *Environ. Res. Lett.* 11. doi:10.1088/1748-9326/11/2/024015
- European Commission, 2016. Monitoring Agricultural Resources (MARS) [WWW Document]. URL <https://ec.europa.eu/jrc/en/mars> (accessed 4.9.17).
- FAO, 2014. SAFA Sustainability Assessment of Food and Agriculture Systems: Guidelines (Version 3.0). Rome.
- FAO, 2013. SAFA (Sustainability Assessment of Food and Agriculture Systems) Indicators.
- FAO, 2009. How to Feed the World in 2050. http://www.fao.org/fileadmin/templates/wsfs/docs/expert_paper/How_to_Feed_the_World_in_2050.pdf.
- FAO, 2004. The ethics of sustainable agricultural intensification, FAO Ethics Series 3. <ftp://ftp.fao.org/docrep/fao/007/j0902e/j0902e00.pdf>.
- Farrell, A., Hart, M., 1998. What does sustainability really mean?: The search for useful indicators. *Environ. Sci. Policy Sustain. Dev.* 40, 4–31. doi:10.1080/00139159809605096
- Feng, X., Fu, B., Yang, X., Lü, Y., 2010. Remote sensing of ecosystem services: An opportunity for spatially explicit assessment. *Chinese Geogr. Sci.* 20, 522–535. doi:10.1007/s11769-010-0428-y
- Ferencz, C., Bognár, P., Lichtenberger, J., Hamar, D., Tarcsai, G., Timár, G., Molnár, G., Pásztor, S.Z., Steinbach, P., Székely, B., Ferencz, O.E., Ferencz-Árkos, I., 2004. Crop yield estimation by satellite remote sensing. *Int. J. Remote Sens.* 25, 4113–4149. doi:10.1080/01431160410001698870
- Firbank, L.G., Elliott, J., Drake, B., Cao, Y., Gooday, R., 2013. Evidence of sustainable intensification among British farms. *Agric. Ecosyst. Environ.* 173, 58–65. doi:10.1016/j.agee.2013.04.010
- Foley, J.A., Ruth DeFries, Gregory P. Asner, Carol Barford, Gordon Bonan, Stephen R. Carpenter, F. Stuart Chapin,

- Michael T. Coe, Gretchen C. Daily, Holly K. Gibbs, Joseph H. Helkowski, Tracey Holloway, Erica A. Howard, Christopher J. Kucharik, Chad Monfreda, Jonathan A. Patz, I. Colin Prentice, Navin Ramankutty, Snyder, P.K., 2005. Global consequences of land use. *Science* (80-.). 309, 570–574.
- Foody, G.M., 2004. Spatial nonstationarity and scale-dependency in the relationship between species richness and environmental determinants for the sub-Saharan endemic avifauna. *Glob. Ecol. Biogeogr.* 13, 315–320. doi:10.1111/j.1466-822X.2004.00097.x
- Galli, A., Wiedmann, T., Ercin, E., Knoblauch, D., Ewing, B., Giljum, S., 2012. Integrating Ecological, Carbon and Water footprint into a “footprint Family” of indicators: Definition and role in tracking human pressure on the planet. *Ecol. Indic.* 16, 100–112. doi:10.1016/j.ecolind.2011.06.017
- Garnett, T., Appleby, M.C., Balmford, A., Bateman, I.J., Benton, T.G., Bloomer, P., Burlingame, B., Dawkins, M., Dolan, L., Fraser, D., Herrero, M., Hoffman, I., Smith, P., Thornton, P.K., Toulmin, C., Vermeulen, S.J., Godfray, H.C.J., 2013. Sustainable intensification in agriculture: premises and policies. *Science* (80-.). 341, 33–34. doi:10.1126/science.1234485
- Garnett, T., Godfray, H.C.J., 2012. Sustainable intensification in agriculture: Navigating a course through competing food system priorities. A report on a workshop. Oxford, UK: Food Climate Research Network and the Oxford Martin Programme on the Future of Food, University of Oxford.
- GEO, n.d. GEOGLAM [WWW Document]. URL <http://www.geoglam.org/index.php/en/> (accessed 4.9.17).
- Gevaert, C.M., García-Haro, F.J., 2015. A comparison of STARFM and an unmixing-based algorithm for Landsat and MODIS data fusion. *Remote Sens. Environ.* 156, 34–44. doi:10.1016/j.rse.2014.09.012
- Gillespie, T.W., Foody, G.M., Rocchini, D., Giorgi, A.P., Saatchi, S., 2008. Measuring and modelling biodiversity from space. *Prog. Phys. Geogr.* 32, 203–221. doi:10.1177/0309133308093606
- Godfray, H.C.J., Beddington, J.R., Crute, I.R., Haddad, L., Lawrence, D., Muir, J.F., Pretty, J., Robinson, S., Thomas, S.M., Toulmin, C., 2010. Food Security: The Challenge of Feeding 9 Billion People. *Science* (80-.). 327, 812–818. doi:10.1126/science.1185383
- Godfray, H.C.J., Garnett, T., 2014. Food security and sustainable intensification. *Philos. Trans. R. Soc. B* 369, 2012073. doi:10.1098/rstb.2012.0273
- Gómez-Limón, J.A., Sanchez-Fernandez, G., 2010. Empirical evaluation of agricultural sustainability using composite indicators. *Ecol. Econ.* 69, 1062–1075. doi:10.1016/j.ecolecon.2009.11.027
- Gons, H.J., Auer, M.T., Effler, S.W., 2008. MERIS satellite chlorophyll mapping of oligotrophic and eutrophic waters in the Laurentian Great Lakes. *Remote Sens. Environ.* 112, 4098–4106. doi:10.1016/j.rse.2007.06.029
- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., Moore, R., 2017. Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sens. Environ.* 202, 18–27. doi:10.1016/j.rse.2017.06.031
- Griffiths, G.H., Lee, J., 2000. Landscape pattern and species richness; regional scale analysis from remote sensing. *Int. J. Remote Sens.* 21, 2685–2704. doi:10.1080/01431160050110232
- Griffiths, P., van der Linden, S., Kuemmerle, T., Hostert, P., 2013. A Pixel-Based Landsat Compositing Algorithm for Large Area Land Cover Mapping. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* 6, 2088–2101. doi:10.1109/JSTARS.2012.2228167
- Gustafson, E.J., 1998. Quantifying Landscape Spatial Pattern: What Is the State of the Art? *Ecosystems* 1, 143–156. doi:10.1007/s100219900011
- Häni, F., Braga, F., Stämpfli, A., Keller, T., Fischer, M., Porsche, H., 2003. RISE, a tool for holistic sustainability assessment at the farm level. *Int. Food Agribus. Manag. Rev.* 6, 78–90.
- Häni, F.J., Stämpfli, A., Gerber, T., Porsche, H., Thalmann, C., Studer, C., 2006. RISE: A tool for improving sustainability

- in agriculture - a case study with tea farms in southern India, in: Häni, F.J., Pintér, L., Herren, H.R. (Eds.), *Sustainable Agriculture: From Principles to Common Practice. Proceedings and Outputs of the First Symposium of the International Forum on Assessing Sustainability in Agriculture (INFASA)*, March 16, 2006, Bern, Switzerland. International Institute for Sustainable Development, pp. 121–148.
- Hein, L., 2014. Biophysical Modelling and Analysis of Ecosystem Services in an Ecosystem Accounting Context DRAFT. <http://img.teebweb.org/wp-content/uploads/2017/01/ANCA-Tech-Guid-9.pdf>.
- Hermosilla, T., Wulder, M.A., White, J.C., Coops, N.C., Hobart, G.W., 2018. Disturbance-Informed Annual Land Cover Classification Maps of Canada's Forested Ecosystems for a 29-Year Landsat Time Series. *Can. J. Remote Sens.* 44, 67–87. doi:10.1080/07038992.2018.1437719
- Herzog, F., Steiner, B., Bailey, D., Baudry, J., Billeter, R., Bukáček, R., De Blust, G., De Cock, R., Dirksen, J., Dormann, C.F., De Filippi, R., Frossard, E., Liira, J., Schmidt, T., Stöckli, R., Thenail, C., Van Wingerden, W., Bugter, R., 2006. Assessing the intensity of temperate European agriculture at the landscape scale. *Eur. J. Agron.* 24, 165–181. doi:10.1016/j.eja.2005.07.006
- Hollmann, R., Merchant, C.J., Saunders, R., Downy, C., Buchwitz, M., Cazenave, A., Chuvieco, E., Defourny, P., de Leeuw, G., Forsberg, R., Holzer-Popp, T., Paul, F., Sandven, S., Sathyendranath, S., van Roozendaal, M., Wagner, W., 2013. The ESA Climate Change Initiative: Satellite Data Records for Essential Climate Variables. *Bull. Am. Meteorol. Soc.* 94, 1541–1552. doi:10.1175/bams-d-11-00254.1
- Honnay, O., Piessens, K., Van Landuyt, W., Hermy, M., Gulinck, H., 2003. Satellite based land use and landscape complexity indices as predictors for regional plant species diversity. *Landsc. Urban Plan.* 63, 241–250. doi:10.1016/S0169-2046(02)00194-9
- Inglada, J., Vincent, A., Arias, M., Tardy, B., Morin, D., Rodes, I., 2017. Operational High Resolution Land Cover Map Production at the Country Scale Using Satellite Image Time Series. *Remote Sens.* 9, 95. doi:10.3390/rs9010095
- Jackson, T.J., Chen, D., Cosh, M., Li, F., Anderson, M., Walthall, C., Doriaswamy, P., Hunt, E.R., 2004. Vegetation water content mapping using Landsat data derived normalized difference water index for corn and soybeans. *Remote Sens. Environ.* 92, 475–482. doi:10.1016/j.rse.2003.10.021
- Kandasamy, S., Baret, F., Verger, A., Neveux, P., Weiss, M., 2013. A comparison of methods for smoothing and gap filling time series of remote sensing observations - application to MODIS LAI products. *Biogeosciences* 10, 4055–4071. doi:10.5194/bg-10-4055-2013
- Kerr, J.T., Cihlar, J., 2003. Land use and cover with intensity of agriculture for Canada from satellite and census data. *Glob. Ecol. Biogeogr.* 12, 161–172. doi:10.1046/j.1466-822X.2003.00017.x
- Kerr, J.T., Ostrovsky, M., 2003. From space to species: Ecological applications for remote sensing. *Trends Ecol. Evol.* 18, 299–305. doi:10.1016/S0169-5347(03)00071-5
- Kissling, W.D., Ahumada, J.A., Bowser, A., Fernandez, M., Fernández, N., García, E.A., Guralnick, R.P., Isaac, N.J.B., Kelling, S., Los, W., McRae, L., Mihoub, J.B., Obst, M., Santamaria, M., Skidmore, A.K., Williams, K.J., Agosti, D., Amariles, D., Arvanitidis, C., Bastin, L., De Leo, F., Egloff, W., Elith, J., Hobern, D., Martin, D., Pereira, H.M., Pesole, G., Peterseil, J., Saarenmaa, H., Schigel, D., Schmeller, D.S., Segata, N., Turak, E., Uhler, P.F., Wee, B., Hardisty, A.R., 2018. Building essential biodiversity variables (EBVs) of species distribution and abundance at a global scale. *Biol. Rev.* 93, 600–625. doi:10.1111/brev.12359
- Kuemmerle, T., Erb, K., Meyfroidt, P., Müller, D., Verburg, P.H., Estel, S., Haberl, H., Hostert, P., Jepsen, M.R., Kastner, T., Levers, C., Lindner, M., Plutzer, C., Verkerk, P.J., van der Zanden, E.H., Reenberg, A., 2013. Challenges and opportunities in mapping land use intensity globally. *Curr. Opin. Environ. Sustain.* 5, 484–493. doi:10.1016/j.cosust.2013.06.002
- Kuenzer, C., Ottinger, M., Wegmann, M., Guo, H., Wang, C., Zhang, J., Dech, S., Wikelski, M., 2014. Earth observation satellite sensors for biodiversity monitoring: potentials and bottlenecks. *Int. J. Remote Sens.* 35, 6599–6647. doi:10.1080/01431161.2014.964349

- Lakhankar, T., Ghedira, H., Temimi, M., Azar, A.E., Khanbilvardi, R., 2009. Effect of land cover heterogeneity on soil moisture retrieval using active microwave remote sensing data. *Remote Sens.* 1, 80–91. doi:10.3390/rs1020080
- Lampkin, N.H., Pearce, B.D., Leake, A.R., Creissen, H., Gerrard, C.L., Girling, R., Lloyd, S., Padel, S., Smith, J., Smith, L.G., Vieweger, A., Wolfe, M.S., 2015. The Role of Agroecology in Sustainable Intensification. Report for the Land Use Policy Group. Organic Research Centre, Elm Farm and Game & Wildlife Conservation Trust.
- Lasne, Y., Paillou, P., Ruffié, G., Serradilla, C., Freeman, A., Farr, T., McDonald, K., Chapman, B., 2008. Effect of Salinity on the Dielectric Properties of Geological Materials: Implication for Soil Moisture Detection by Means of Remote Sensing. *IEEE Trans. Geosci. Remote Sens.* 6, 1674–1688. doi:10.1109/TGRS.2008.916220
- Li, J., Roy, D.P., 2017. A global analysis of Sentinel-2A, Sentinel-2B and Landsat-8 data revisit intervals and implications for terrestrial monitoring. *Remote Sens.* 9, 902. doi:10.3390/rs9090902
- Li, Z., Guo, X., 2012. Detecting climate effects on vegetation in northern mixed rairie using NOAA AVHRR 1-km time-series NDVI data. *Remote Sens.* 4, 120–134. doi:10.3390/rs4010120
- Li, Z., Xu, D., Guo, X., 2014. Remote sensing of ecosystem health: opportunities, challenges, and future perspectives. *Sensors* 14, 21117–21139. doi:10.3390/s141121117
- Loveland, T.R., Reed, B.C., Ohlen, D.O., Brown, J.F., Zhu, Z., Yang, L., Merchant, J.W., 2000. Development of a global land cover characteristics database and IGBP DISCover from 1 km AVHRR data. *Int. J. Remote Sens.* 21, 1303–1330. doi:10.1080/014311600210191
- Luoto, M., Toivonen, T., Heikkinen, R.K., 2002. Prediction of total and rare plant species richness in agricultural landscapes from satellite images and topographic data. *Landsc. Ecol.* 17, 195–217. doi:10.1023/A:1020288509837
- Mairota, P., Cafarelli, B., Didham, R.K., Lovergine, F.P., Lucas, R.M., Nagendra, H., Rocchini, D., Tarantino, C., 2015. Challenges and opportunities in harnessing satellite remote-sensing for biodiversity monitoring. *Ecol. Inform.* 30, 207–214. doi:10.1016/j.ecoinf.2015.08.006
- Martinez, J.M., Guyot, J.L., Cochonneau, G., 2008. Monitoring of surface water quality in large rivers with satellite imagery - Application to the Amazon basin. www.iwra.org/congress/resource/abs649_article.pdf.
- Matson, P.A., Parton, W.J., Power, A.G., Swift, M.J., 1997. Agricultural intensification and ecosystem properties. *Science* (80-.). 277, 504–509. doi:10.1126/science.277.5325.504
- Meeus, J.H.A., 1993. The transformation of agricultural landscapes in Western Europe. *Sci. Total Environ.* 129, 171–190. doi:10.1016/0048-9697(93)90169-7
- Mingwei, Z., Qingbo, Z., Zhongxin, C., Jia, L., Yong, Z., Chongfa, C., 2008. Crop discrimination in Northern China with double cropping systems using Fourier analysis of time-series MODIS data. *Int. J. Appl. Earth Obs. Geoinf.* 10, 476–485. doi:10.1016/j.jag.2007.11.002
- Mitchell, G., May, A., McDonald, A., 1995. PICABUE: a methodological framework for the development of indicators of sustainable development. *Int. J. Sustain. Dev. World Ecol.* 2, 104–123. doi:10.1080/13504509509469893
- Moriondo, M., Maselli, F., Bindi, M., 2007. A simple model of regional wheat yield based on NDVI data. *Eur. J. Agron.* 26, 266–274. doi:10.1016/j.eja.2006.10.007
- Mueller, N., Lewis, A., Roberts, D., Ring, S., Melrose, R., Sixsmith, J., Lymburner, L., McIntyre, A., Tan, P., Curnow, S., Ip, A., 2016. Water observations from space: Mapping surface water from 25 years of Landsat imagery across Australia. *Remote Sens. Environ.* 174, 341–352. doi:10.1016/j.rse.2015.11.003
- Musumba, M., Palm, C., Grabowski, P., Snapp, S.S., 2017. A Framework for Selecting and Analyzing Indicators of Sustainable intensification.
- Myneni, R.B., Nemani, R.R., Running, S.W., 1997. Estimation of global leaf area index and absorbed PAR using

- radiative transfer models. *IEEE Trans. Geosci. Remote Sens.* 35, 1380–1393. doi:10.1109/36.649788
- Nagendra, H., 2001. Using remote sensing to assess biodiversity. *Int. J. Remote Sens.* 22, 2377–2400. doi:10.1080/01431160117096
- Newton, I., 2004. The recent declines of farmland bird populations in Britain: An appraisal of causal factors and conservation actions. *Ibis (Lond. 1859)*. 146, 579–600. doi:10.1111/j.1474-919X.2004.00375.x
- Niedertscheider, M., Kastner, T., Fetzl, T., Haberl, H., Kroisleitner, C., Plutzer, C., Erb, K.-H., 2016. Mapping and analysing cropland use intensity from a NPP perspective. *Environ. Res. Lett.* 11. doi:10.1088/1748-9326/11/1/014008
- OneSoil, 2018. OneSoil [WWW Document]. URL <https://onesoil.ai/en/> (accessed 4.8.19).
- Pacheco, A., McNairn, H., 2010. Evaluating multispectral remote sensing and spectral unmixing analysis for crop residue mapping. *Remote Sens. Environ.* 114, 2219–2228. doi:10.1016/j.rse.2010.04.024
- Pereira, H.M., Ferrier, S., Walters, M., Geller, G.N., Jongman, R.H.G., Scholes, R.J., Bruford, M.W., Brummitt, N., Butchart, S.H.M., Cardoso, A.C., Coops, N.C., Dulloo, E., Faith, D.P., Freyhof, J., Gregory, R.D., Heip, C., Höft, R., Hurtt, G., Jetz, W., Karp, D.S., McGeoch, M.A., Obura, D., Onoda, Y., Pettorelli, N., Reyers, B., Sayre, R., Scharlemann, J.P.W., Stuart, S.N., Turak, E., Walpole, M., Wegmann, M., 2013. Essential biodiversity variables. *Science (80-.)*. 339, 277–278. doi:10.1126/science.1229931
- Pettorelli, N., Laurance, W.F., O'Brien, T.G., Wegmann, M., Nagendra, H., Turner, W., 2014. Satellite remote sensing for applied ecologists: Opportunities and challenges. *J. Appl. Ecol.* 51, 839–848. doi:10.1111/1365-2664.12261
- Pettorelli, N., Wegmann, M., Skidmore, A., Mùcher, S., Dawson, T.P., Fernandez, M., Lucas, R., Schaepman, M.E., Wang, T., O'Connor, B., Jongman, R.H.G., Kempeneers, P., Sonnenschein, R., Leidner, A.K., Böhm, M., He, K.S., Nagendra, H., Dubois, G., Fatoyinbo, T., Hansen, M.C., Paganini, M., de Klerk, H.M., Asner, G.P., Kerr, J.T., Estes, A.B., Schmeller, D.S., Heiden, U., Rocchini, D., Pereira, H.M., Turak, E., Fernandez, N., Lausch, A., Cho, M.A., Alcaraz-Segura, D., McGeoch, M.A., Turner, W., Mueller, A., St-Louis, V., Penner, J., Vihervaara, P., Belward, A., Reyers, B., Geller, G.N., 2016. Framing the concept of satellite remote sensing essential biodiversity variables: challenges and future directions. *Remote Sens. Ecol. Conserv.* 2, 122–131. doi:10.1002/rse2.15
- Power, A.G., 2010. Ecosystem services and agriculture: tradeoffs and synergies. *Philos. Trans. R. Soc. B* 365, 2959–2971. doi:10.1098/rstb.2010.0143
- Prasad, A.K., Chai, L., Singh, R.P., Kafatos, M., 2006. Crop yield estimation model for Iowa using remote sensing and surface parameters. *Int. J. Appl. Earth Obs. Geoinf.* 8, 26–33. doi:10.1016/j.jag.2005.06.002
- Pretty, J., 2008. Agricultural sustainability: concepts, principles and evidence. *Philos. Trans. R. Soc. B* 363, 447–465. doi:10.1098/rstb.2007.2163
- Pretty, J., Bharucha, Z.P., 2014. Sustainable intensification in agricultural systems. *Ann. Bot.* 114, 1571–1596. doi:10.1093/aob/mcu205
- Pretty, J., Toulmin, C., Williams, S., 2011. Sustainable intensification in African agriculture. *Int. J. Agric. Sustain.* 9, 5–24. doi:10.3763/ijas.2010.0583
- Rasul, G., Thapa, G.B., 2004. Sustainability of ecological and conventional agricultural systems in Bangladesh: An assessment based on environmental, economic and social perspectives. *Agric. Syst.* 79, 327–351. doi:10.1016/S0308-521X(03)00090-8
- Rigby, D., Woodhouse, P., Young, T., Burton, M., 2001. Constructing a farm level indicator of sustainable agricultural practice. *Ecol. Econ.* 39, 463–478. doi:10.1016/S0921-8009(01)00245-2
- Rocchini, D., Balkenhol, N., Carter, G.A., Foody, G.M., Gillespie, T.W., He, K.S., Kark, S., Levin, N., Lucas, K., Luoto, M., Nagendra, H., Oldeland, J., Ricotta, C., Southworth, J., Neteler, M., 2010. Remotely sensed spectral heterogeneity as a proxy of species diversity: Recent advances and open challenges. *Ecol. Inform.* 5, 318–329.

- Rodrigues, G.S., Rodrigues, I.A., Buschinelli, C.C. de A., de Barros, I., 2010. Integrated farm sustainability assessment for the environmental management of rural activities. *Environ. Impact Assess. Rev.* 30, 229–239. doi:10.1016/j.eiar.2009.10.002
- Roy, D.P., Ju, J., Kline, K., Scaramuzza, P.L., Kovalsky, V., Hansen, M., Loveland, T.R., Vermote, E., Zhang, C., 2010. Web-enabled Landsat Data (WELD): Landsat ETM+ composited mosaics of the conterminous United States. *Remote Sens. Environ.* 114, 35–49. doi:10.1016/j.rse.2009.08.011
- Roy, R., Chan, N.W., 2012. An assessment of agricultural sustainability indicators in Bangladesh: Review and synthesis. *Environmentalist* 32, 99–110. doi:10.1007/s10669-011-9364-3
- Schut, M., van Asten, P., Okafor, C., Hicintuka, C., Mapatano, S., Nabahungu, N.L., Kagabo, D., Muchunguzi, P., Njukwe, E., Dongsop-Nguezet, P.M., Sartas, M., Vanlauwe, B., 2016. Sustainable intensification of agricultural systems in the Central African Highlands: The need for insitutional innovation. *Agric. Syst.* 145, 165–176. doi:10.1016/j.agsy.2016.03.005
- Senf, C., Leitão, P.J., Pflugmacher, D., van der Linden, S., Hostert, P., 2015. Mapping land cover in complex Mediterranean landscapes using Landsat: Improved classification accuracies from integrating multi-seasonal and synthetic imagery. *Remote Sens. Environ.* 156, 527–536. doi:10.1016/j.rse.2014.10.018
- Shao, Y., Lunetta, R.S., Wheeler, B., Iiames, J.S., Campbell, J.B., 2016. An evaluation of time-series smoothing algorithms for land-cover classifications using MODIS-NDVI multi-temporal data. *Remote Sens. Environ.* 174, 258–265. doi:10.1016/j.rse.2015.12.023
- Shoshany, M., Goldshleger, N., Chudnovsky, A., 2013. Monitoring of agricultural soil degradation by remote-sensing methods: a review. *Int. J. Remote Sens.* 34, 6152–6181. doi:10.1080/01431161.2013.793872
- Shriar, A.J., 2000. Agricultural intensity and its measurement in frontier regions. *Agrofor. Syst.* 49, 301–318. doi:10.1023/A:1006316131781
- Singh, R., Semwal, D.P., Rai, A., Chhikara, R.S., 2002. Small area estimation of crop yield using remote sensing satellite data. *Int. J. Remote Sens.* 23, 49–56. doi:10.1080/01431160010014756
- Singh, R.K., Murty, H.R., Gupta, S.K., Dikshit, A.K., 2009. An overview of sustainability assessment methodologies. *Ecol. Indic.* 9, 189–212. doi:10.1016/j.ecolind.2008.05.011
- Smith, A., Snapp, S., Chikowo, R., Thorne, P., Bekunda, M., Glover, J., 2017. Measuring sustainable intensification in smallholder agroecosystems: A review. *Glob. Food Sec.* 12, 127–138. doi:10.1016/j.gfs.2016.11.002
- Snapp, S.S., Grabowski, P., Chikowo, R., Smith, A., Anders, E., Sirrine, D., Chimonyo, V., Bekunda, M., 2018. Maize yield and profitability tradeoffs with social, human and environmental performance: Is sustainable intensification feasible? *Agric. Syst.* 162, 77–88. doi:10.1016/j.agsy.2018.01.012
- Stein, A., Riley, J., Halberg, N., 2001. Issues of scale for environmental indicators. *Agric. Ecosyst. Environ.* 87, 215–232. doi:10.1016/S0167-8809(01)00280-8
- Stoate, C., Boatman, N.D., Borralho, R.J., Carvalho, C.R., de Snoo, G.R., Eden, P., 2001. Ecological impacts of arable intensification in Europe. *J. Environ. Manage.* 63, 337–365. doi:10.1006/jema.2001.0473
- Sulistioadi, Y.B., Tseng, K.H., Shum, C.K., Hidayat, H., Sumaryono, M., Suhardiman, A., Setiawan, F., Sunarso, S., 2015. Satellite radar altimetry for monitoring small rivers and lakes in Indonesia. *Hydrol. Earth Syst. Sci.* 19, 341–359. doi:10.5194/hess-19-341-2015
- Tan, B., Morissette, J.T., Wolfe, R.E., Gao, F., Ederer, G.A., Nightingale, J., Pedelty, J.A., 2011. An Enhanced TIMESAT Algorithm for Estimating Vegetation Phenology Metrics From MODIS Data. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* 4, 361–371. doi:10.1109/JSTARS.2010.2075916

- Tebbs, E., Rowland, C., Smart, S., Maskell, L., Norton, L., 2017. Regional-Scale High Spatial Resolution Mapping of Aboveground Net Primary Productivity (ANPP) from Field Survey and Landsat Data: A Case Study for the Country of Wales. *Remote Sens.* 9, 801. doi:10.3390/rs9080801
- Teillard, F., Allaire, G., Cahuzac, E., Léger, F., Maigné, E., Tichit, M., 2012. A novel method for mapping agricultural intensity reveals its spatial aggregation: Implications for conservation policies. *Agric. Ecosyst. Environ.* 149, 135–143. doi:10.1016/j.agee.2011.12.018
- Teluguntla, P., Thenkabail, P., Oliphant, A., Xiong, J., Gumma, M.K., Congalton, R.G., Yadav, K., Huete, A., 2018. A 30-m landsat-derived cropland extent product of Australia and China using random forest machine learning algorithm on Google Earth Engine cloud computing platform. *ISPRS J. Photogramm. Remote Sens.* 144, 325–340. doi:10.1016/j.isprsjprs.2018.07.017
- Temme, A.J.A.M., Verburg, P.H., 2011. Mapping and modelling of changes in agricultural intensity in Europe. *Agric. Ecosyst. Environ.* 140, 46–56. doi:10.1016/j.agee.2010.11.010
- Tilman, D., Balzer, C., Hill, J., Befort, B.L., 2011. Global food demand and the sustainable intensification of agriculture. *PNAS* 108, 20260–20264. doi:10.1073/pnas.1116437108
- Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R., Polasky, S., 2002. Agricultural sustainability and intensive production practices. *Nature* 418, 671–677. doi:10.1038/nature01014
- Torres, R., Snoeij, P., Geudtner, D., Bibby, D., Davidson, M., Attema, E., Potin, P., Rommen, B.Ö., Floury, N., Brown, M., Traver, I.N., Deghaye, P., Duesmann, B., Rosich, B., Miranda, N., Bruno, C., L'Abbate, M., Croci, R., Pietropaolo, A., Huchler, M., Rostan, F., 2012. GMES Sentinel-1 mission. *Remote Sens. Environ.* 120, 9–24. doi:10.1016/j.rse.2011.05.028
- Trombetti, M., Riaño, D., Rubio, M.A., Cheng, Y.B., Ustin, S.L., 2008. Multi-temporal vegetation canopy water content retrieval and interpretation using artificial neural networks for the continental USA. *Remote Sens. Environ.* 112, 203–215. doi:10.1016/j.rse.2007.04.013
- Turner, W., Rondinini, C., Pettorelli, N., Mora, B., Leidner, A.K., Szantoi, Z., Buchanan, G., Dech, S., Dwyer, J., Herold, M., Koh, L.P., Leimgruber, P., Taubenboeck, H., Wegmann, M., Wikelski, M., Woodcock, C., 2015. Free and open-access satellite data are key to biodiversity conservation. *Biol. Conserv.* 182, 173–176. doi:10.1016/j.biocon.2014.11.048
- Turner, W., Spector, S., Gardiner, N., Fladeland, M., Sterling, E., Steininger, M., 2003. Remote sensing for biodiversity science and conservation. *Trends Ecol. Evol.* 18, 306–314. doi:10.1016/S0169-5347(03)00070-3
- USDA FAS, n.d. GLAM - Global Agricultural Monitoring [WWW Document]. URL <https://www.pecad.fas.usda.gov/glam.htm> (accessed 4.9.17).
- Van Cauwenbergh, N., Biala, K., Bielders, C., Brouckaert, V., Franchois, L., Garcia Ciudad, V., Hermy, M., Mathijs, E., Muys, B., Reijnders, J., Sauvenier, X., Valckx, J., Vanclooster, M., Van der Veken, B., Wauters, E., Peeters, A., 2007. SAFE-A hierarchical framework for assessing the sustainability of agricultural systems. *Agric. Ecosyst. Environ.* 120, 229–242. doi:10.1016/j.agee.2006.09.006
- Van Passel, S., Meul, M., 2012. Multilevel and multi-user sustainability assessment of farming systems. *Environ. Impact Assess. Rev.* 32, 170–180. doi:10.1016/j.eiar.2011.08.005
- Verger, A., Baret, F., Weiss, M., Kandasamy, S., Vermote, E., 2013. The CACAO method for smoothing, gap filling, and characterizing seasonal anomalies in satellite time series. *IEEE Trans. Geosci. Remote Sens.* 51, 1963–1972. doi:10.1109/TGRS.2012.2228653
- Vuolo, F., Ng, W.-T., Atzberger, C., 2017. Smoothing and gap-filling of high resolution multi-spectral time series: Example of Landsat data. *Int. J. Appl. Earth Obs. Geoinf.* 57, 202–213. doi:10.1016/j.jag.2016.12.012
- Wagner, W., Naeimi, V., Scipal, K., Jeu, R., Martínez-Fernández, J., 2007. Soil moisture from operational meteorological satellites. *Hydrogeol. J.* 15, 121–131. doi:10.1007/s10040-006-0104-6


- Wang, K., Franklin, S.E., Guo, X., Cattet, M., 2010. Remote sensing of ecology, biodiversity and conservation: A review from the perspective of remote sensing specialists. *Sensors* 10, 9647–9667. doi:10.3390/s101109647
- Wang, Q., Atkinson, P.M., 2018. Spatio-temporal fusion for daily Sentinel-2 images. *Remote Sens. Environ.* 204, 31–42. doi:10.1016/j.rse.2017.10.046
- Weiers, S., Bock, M., Wissen, M., Rossner, G., 2004. Mapping and indicator approaches for the assessment of habitats at different scales using remote sensing and GIS methods. *Landsc. Urban Plan.* 67, 43–65. doi:10.1016/S0169-2046(03)00028-8
- Whitcraft, A.K., Becker-Reshef, I., Justice, C.O., 2015. A framework for defining spatially explicit earth observation requirements for a global agricultural monitoring initiative (GEOGLAM). *Remote Sens.* 7, 1461–1481. doi:10.3390/rs70201461
- Wiek, A., Binder, C., 2005. Solution spaces for decision-making - A sustainability assessment tool for city-regions. *Environ. Impact Assess. Rev.* 25, 589–608. doi:10.1016/j.eiar.2004.09.009
- Xiong, J., Thenkabail, P.S., Tilton, J.C., Gumma, M.K., Teluguntla, P., Oliphant, A., Congalton, R.G., Yadav, K., Gorelick, N., 2017. Nominal 30-m cropland extent map of continental Africa by integrating pixel-based and object-based algorithms using Sentinel-2 and Landsat-8 data on google earth engine. *Remote Sens.* 9, 1065. doi:10.3390/rs9101065
- Yan, H., Xiao, X., Huang, H., Liu, J., Chen, J., Bai, X., 2014. Multiple Cropping Intensity in China Derived from Agro-meteorological Observations and MODIS Data. *Chinese Geogr. Sci.* 24, 205–219. doi:10.1007/s11769-013-0637-2
- Zahm, F., Viaux, P., Vilain, L., Girardin, P., Mouchet, C., 2008. Assessing Farm Sustainability with the IDEA Method - from the Concept of Agriculture Sustainability to Case Studies on Farms. *Sustain. Dev.* 16. doi:10.1002/sd.380
- Zhang, Y., Atkinson, P.M., Li, X., Ling, F., Wang, Q., Du, Y., 2017. Learning-Based Spatial-Temporal Superresolution Mapping of Forest Cover with MODIS Images. *IEEE Trans. Geosci. Remote Sens.* 55, 600–614. doi:10.1109/TGRS.2016.2613140
- Zhen, L., Routray, J.K., Zoebisch, M.A., Chen, G., Xie, G., Cheng, S., 2005. Three dimensions of sustainability of farming practices in the North China Plain: A case study from Ningjin County of Shandong Province, PR China. *Agric. Ecosyst. Environ.* 105, 507–522. doi:10.1016/j.agee.2004.07.012

Supplementary material

Contents

Page number	Table/figure number	Description
2	Figure S1	<i>Some of the key environmental impacts of various mechanisms of agricultural intensification with references. Reference numbers correspond to the list on pages 3-4.</i>
3-4		<i>References for figure S1</i>
5-6	Table S1	<i>Examples of data sources and indicators used by various authors to investigate agricultural intensity.</i>
7	Table S2	<i>Example EO-based methods used by researchers to assess agricultural intensity.</i>
8	Table S3	<i>Example EO-based methods used by researchers to assess vegetation health.</i>
9-10	Table S4	<i>Example EO-based methods used by researchers to assess soil quality and soil erosion/protection.</i>
11	Table S5	<i>Example EO-based methods used by researchers to assess water quality and water availability.</i>
12	Table S6	<i>Example EO-based methods used by researchers to assess biodiversity.</i>
13	Table S7	<i>Example EO-based methods used by researchers to assess ecosystem health.</i>
14-25		<i>References for tables S1-S7</i>

Figure S1: Some of the key environmental impacts of various mechanisms of agricultural intensification with references. Reference numbers correspond to the list on pages 3-4.



References for figure S1

1. Alauddin, M., Quiggin, J., 2008. Agricultural intensification, irrigation and the environment in South Asia: Issues and policy options. *Ecol. Econ.* 65, 111–124. doi:10.1016/j.ecolecon.2007.06.004
2. Baum, K.A., Haynes, K.J., Dilleuth, F.P., Cronin, J.T., 2004. The matrix enhances the effectiveness of corridors and stepping stones. *Reports Ecol.* 85, 2671–2676. doi:10.1890/04-0500
3. Benckiser, G., 1997. Organic inputs and soil metabolism, in: Benckiser, G. (Ed.), *Fauna in Soil Ecosystems: Recycling Processes, Nutrient Fluxes and Agricultural Production*. Dekker, New York, pp. 7–62.
4. Benton, T.G., Vickery, J.A., Wilson, J.D., 2003. Farmland biodiversity: Is habitat heterogeneity the key? *Trends Ecol. Evol.* 18, 182–188. doi:10.1016/S0169-5347(03)00011-9
5. Blažek, P., Lepš, J., 2015. Victims of agricultural intensification: Mowing date affects *Rhinanthus* spp. regeneration and fruit ripening. *Agric. Ecosyst. Environ.* 211, 10–16. doi:10.1016/j.agee.2015.04.022
6. Campbell, L.H., Avery, M.I., Donald, P., Evans, A.D., Green, R.E., Wilson, J.D., 1997. A review of the indirect effects of pesticides on birds. JNCC Report No. 277. Peterborough.
7. Ciccolini, V., Bonari, E., Ercoli, L., Pellegrino, E., 2016. Phylogenetic and multivariate analyses to determine the effect of agricultural land-use intensification and soil physico-chemical properties on N-cycling microbial communities in drained Mediterranean peaty soils. *Biol. Fertil. Soils* 52, 811–824. doi:10.1007/s00374-016-1121-9
8. Crowder, D.W., Jabbour, R., 2014. Relationships between biodiversity and biological control in agroecosystems: Current status and future challenges. *Biol. Control* 75, 8–17. doi:10.1016/j.biocontrol.2013.10.010
9. De Jong, S.M., Paracchini, M.L., Bertolo, F., Folving, S., Megier, J., De Roo, A.P.J., 1999. Regional assessment of soil erosion using the distributed model SEMMED and remotely sensed data. *Catena* 37, 291–308. doi:10.1016/S0341-8162(99)00038-7
10. Donald, P.F., Green, R.E., Heath, M.F., 2001. Agricultural intensification and the collapse of Europe's farmland bird populations. *Proc. R. Soc. B* 268, 25–29. doi:10.1098/rspb.2000.1325
11. Dorrough, J., Scroggie, M.P., 2008. Plant responses to agricultural intensification. *J. Appl. Ecol.* 45, 1274–1283. doi:10.1111/j.1365-2664.2008.01501.x
12. Douglas, E.M., Beltrán-Przekurat, A., Niyogi, D., Pielke, R.A., Vörösmarty, C.J., 2009. The impact of agricultural intensification and irrigation on land-atmosphere interactions and Indian monsoon precipitation - A mesoscale modeling perspective. *Glob. Planet. Change* 67, 117–128. doi:10.1016/j.gloplacha.2008.12.007
13. Edwards, C.A., 1984. Changes in agricultural practice and their impact on soil organisms, in: Jenkins, D. (Ed.), *Agriculture and the Environment. Proceedings of ITE Symposium no.13 Held at Monks Wood Experimental Station on 28-29 February and 1 March 1984*. pp. 56–65.
14. Firbank, L.G., Petit, S., Smart, S., Blain, A., Fuller, R.J., 2008. Assessing the impacts of agricultural intensification on biodiversity: a British perspective. *Philos. Trans. R. Soc. B Biol. Sci.* 363, 777–787. doi:10.1098/rstb.2007.2183
15. Franke, A.C., Lotz, L.A.P., Van Der Burg, W.J., Van Overbeek, L., 2009. The role of arable weed seeds for agroecosystem functioning. *Weed Res.* 49, 131–141. doi:10.1111/j.1365-3180.2009.00692.x
16. Gabriel, D., Sait, S.M., Hodgson, J.A., Schmutz, U., Kunin, W.E., Benton, T.G., 2010. Scale matters: The impact of organic farming on biodiversity at different spatial scales. *Ecol. Lett.* 13, 858–869. doi:10.1111/j.1461-0248.2010.01481.x
17. Ganasri, B.P., Ramesh, H., 2016. Assessment of soil erosion by RUSLE model using remote sensing and GIS - A case study of Nethravathi Basin. *Geosci. Front.* 7, 953–961. doi:10.1016/j.gsf.2015.10.007
18. González-Estébanez, F.J., García-Tejero, S., Mateo-Tomás, P., Olea, P.P., 2011. Effects of irrigation and landscape heterogeneity on butterfly diversity in Mediterranean farmlands. *Agric. Ecosyst. Environ.* 144, 262–270. doi:10.1016/j.agee.2011.09.002
19. Gordon, L.J., Steffen, W., Jonsson, B.F., Folke, C., Falkenmark, M., Johannessen, A., 2005. Human modification of global water vapor flows from the land surface. *Proc. Natl. Acad. Sci.* 102, 7612–7617. doi:10.1073/pnas.0500208102

20. Guerrero, I., Morales, M.B., Oñate, J.J., Geiger, F., Berendse, F., Snoo, G. de, Eggers, S., Pärt, T., Bengtsson, J., Clement, L.W., Weisser, W.W., Olszewski, A., Ceryngier, P., Hawro, V., Liira, J., Aavik, T., Fischer, C., Flohre, A., Thies, C., Tschardtke, T., 2012. Response of ground-nesting farmland birds to agricultural intensification across Europe: Landscape and field level management factors. *Biol. Conserv.* 152, 74–80. doi:10.1016/j.biocon.2012.04.001
21. Herzig, A., Dymond, J., Ausseil, A.-G., 2016. Exploring limits and trade-offs of irrigation and agricultural intensification in the Ruamahanga catchment, New Zealand. *New Zeal. J. Agric. Res.* 59, 216–234. doi:10.1080/00288233.2016.1183685
22. Jeliakov, A., Mimet, A., Chargé, R., Jiguet, F., Devictor, V., Chiron, F., 2016. Impacts of agricultural intensification on bird communities: New insights from a multi-level and multi-facet approach of biodiversity. *Agric. Ecosyst. Environ.* 216, 9–22. doi:10.1016/j.agee.2015.09.017
23. José-María, L., Armengot, L., Blanco-Moreno, J.M., Bassa, M., Sans, F.X., 2010. Effects of agricultural intensification on plant diversity in Mediterranean dryland cereal fields. *J. Appl. Ecol.* 47, 832–840. doi:10.1111/j.1365-2664.2010.01822.x
24. Karp, D.S., Rominger, A.J., Zook, J., Ranganathan, J., Ehrlich, P.R., Daily, G.C., 2012. Intensive agriculture erodes β -diversity at large scales. *Ecol. Lett.* 15, 963–970. doi:10.1111/j.1461-0248.2012.01815.x
25. Kleijn, D., Van Der Voort, L.A.C., 1997. Conservation headlands for rare arable weeds: The effects of fertilizer application and light penetration on plant growth. *Biol. Conserv.* 81, 57–67. doi:10.1016/S0006-3207(96)00153-X
26. Maron, M., Fitzsimons, J.A., 2007. Agricultural intensification and loss of matrix habitat over 23 years in the West Wimmera, south-eastern Australia. *Biol. Conserv.* 135, 587–593. doi:10.1016/j.biocon.2006.10.051
27. Meeus, J.H.A., 1993. The transformation of agricultural landscapes in Western Europe. *Sci. Total Environ.* 129, 171–190.
28. Pielke, R.A., 2001. Influence of the spatial distribution of vegetation and soils on the prediction of cumulus convective rainfall. *Rev. Geophys.* 39, 151–177. doi:10.1029/1999RG000072
29. Roubos, C.R., Rodriguez-Saona, C., Isaacs, R., 2014. Mitigating the effects of insecticides on arthropod biological control at field and landscape scales. *Biol. Control* 75, 28–39. doi:10.1016/j.biocontrol.2014.01.006
30. Stoate, C., Boatman, N.D., Borralho, R.J., Carvalho, C.R., de Snoo, G.R., Eden, P., 2001. Ecological impacts of arable intensification in Europe. *J. Environ. Manage.* 63, 337–365. doi:10.1006/jema.2001.0473
31. Storkey, J., Meyer, S., Still, K.S., Leuschner, C., 2012. The impact of agricultural intensification and land-use change on the European arable flora. *Proc. R. Soc. B Biol. Sci.* 279, 1421–1429. doi:10.1098/rspb.2011.1686
32. Tilman, D., Fargione, J., Wolff, B., D'Antonio, C., Dobson, A., Howarth, R., Schindler, D., Schlesinger, W.H., Simberloff, D., Swackhamer, D., 2001. Forecasting agriculturally driven global environmental change. *Science* (80-.). 292, 281–284. doi:10.1126/science.1057544
33. Tschardtke, T., Klein, A.M., Kruss, A., Steffan-Dewenter, I., Thies, C., 2005. Landscape perspectives on agricultural intensification and biodiversity - Ecosystem service management. *Ecol. Lett.* 8, 857–874. doi:10.1111/j.1461-0248.2005.00782.x
34. Van Oost, K., Govers, G., Desmet, P., 2000. Evaluating the effects of changes in landscape structure on soil erosion by water and tillage. *Landsc. Ecol.* 15, 577–589. doi:10.1023/A:1008198215674
35. Welch, K.D., Harwood, J.D., 2014. Temporal dynamics of natural enemy-pest interactions in a changing environment. *Biol. Control* 75, 18–27. doi:10.1016/j.biocontrol.2014.01.004
36. Williams, P.H., 1982. The distribution and decline of British bumble bees (*Bombus* Latr.). *J. Apic. Res.* 21, 236–245. doi:10.1080/00218839.1982.11100549
37. Wilson, J.D., Evans, J., Browne, S.J., King, J.R., 1997. Territory Distribution and Breeding Success of Skylarks *Alauda arvensis* on Organic and Intensive Farmland in Southern England. *J. Appl. Ecol.* 34, 1462–1478.

1 **Table S1: Examples of data sources and indicators used by various authors to investigate agricultural intensity.**

Study Area	Data sources	Indicator(s)
Beijing mountainous region (Zhang and Li, 2016)	• Statistical Yearbook 2013	• Emergy analysis of agricultural inputs
Northern Spain (Armengot et al., 2011)	• Interviews	• Mean annual inputs of exogenous nitrogen • Weed control intensity • Cereal ratio • Crop diversity • Seed origin
Germany (Egorov et al., 2014)	• Yearly interviews with farmers and land-owners	• Land Use Intensity Index (LUI): summation of fertilization level, mowing frequency and grazing intensity
France (Teillard et al., 2012)	• French FADN (Farm Accountancy Data Network) • Datasets from agricultural social security, CAP declarations, national bovine identification • Topo-climatic data	• Input cost/ha
India (Biradar and Xiao, 2011)	• MODIS (EVI, NDVI, LSWI) • Government agricultural census data • Field ground-truth data inc. crop types and cropping pattern	• Cropping intensity
India (Jain et al., 2013)	• Landsat 5 TM & Landsat 7 ETM+ • MODIS (EVI) • Ground-truth data: Landsat, Quickbird, Worldview & Google Earth Imagery	• Cropping intensity • Multi-cropping
India (Singh et al., 2002)	• Crop cutting experiments • Crop yield estimation surveys • IRS-1B LISS-II	• Crop yield
Northern China (Mingwei et al., 2008)	• MODIS (NDVI)	• Crop acreage
China (Yan et al., 2014)	• Agricultural meteorological stations – crop calendar/crop phenological data • MODIS (EVI) • National LU/LC dataset	• Multi-cropping
China (L. Li et al., 2014)	• MODIS (EVI) • National survey data	• Cropping intensity
China (Xie et al., 2014)	• Secondary agricultural statistics e.g. China Rural Statistical Yearbook	• Emergy analysis of inputs to arable land per ha
United States (Johnson, 2013)	• Cropland Data Layer LC classifications (derived from Landsat TM by Agricultural Statistics Service) • NASS Census of Agriculture & June Acreage Survey	• Area of annually tilled cropland
US Central Great Plains (Wardlow and Egbert, 2008)	• MODIS (NDVI)	• Cropping area
Canada (Kerr and Cihlar, 2003)	• Canadian Census of Agriculture • SPOT 4 VEG	• Agricultural pollution

Table S1 continued

Study area	Data sources	Indicator(s)
Belgium, France, the Netherlands and Switzerland (<i>Le Féon et al., 2010</i>)	• Standardised interviews with farmers	• Global Intensity Index: based on nitrogen input, livestock density and pesticide input
Europe (<i>Donald et al., 2001</i>)	• FAOSTAT database	• PCA analysis based on political & economic differences
Europe (<i>Herzog et al., 2006</i>)	• Interviews • Geo-referenced aerial photographs • European Fourier-Adjusted & Interpolated NDVI dataset (Stockli & Vidale, 2004)	• Nitrogen output • Density of livestock units • Number of pesticide applications
European Union (<i>Reidsma et al., 2006</i>)	• FADN survey	• Input costs • Irrigation use • Organic or not?
European Union (<i>Temme and Verburg, 2011</i>)	• Agricultural statistics & census data • CORINE land cover map • Land Use/Cover Area frame statistical Survey (LUCAS) dataset	• Total nitrogen input
Europe & Turkey (<i>Estel et al., 2016</i>)	• MODIS NDVI (Terra & Aqua satellites) • GlobCorine LC Map • Annual fallow/active crop maps	• Cropping frequency • Multi-cropping • Fallow cycles • Crop duration ratio
Russia (<i>de Beurs and Ioffe, 2014</i>)	• Landsat 5 • MODIS • All-Russia Agricultural Census • Field Observations	• Cropping intensity
Asia (<i>Gray et al., 2014</i>)	• MODIS (EVI)	• Multi-cropping
World (<i>Johnston et al., 2011</i>)	• Global census data from FAOSTAT database – M3 cropland datasets	• Yield gap analysis
World (<i>Niedertscheider et al., 2016</i>)	• Earthstat gridded maps of crop yields & crop area	• Human appropriation of net primary productivity (HANPP)
World (<i>Potter et al., 2010</i>)	• National level fertiliser data – based on questionnaires • Global maps of harvested Area (from Monfreda <i>et al.</i> 2008) • FAO Gridded Livestock of the World maps	• Fertiliser inputs of N & P
World (<i>Siebert et al., 2010</i>)	• MIRCA2000 dataset – monthly growing areas of 26 irrigated & rain-fed crop classes	• Cropping intensity • Crop duration ratio • Extent of fallow land
World (<i>Thenkabail et al., 2009</i>)	• AVHRR • SPOT • JERS-1 • CRU rainfall time series (1961-2000) • Global Elevation dataset • Global Tree Cover data • Google Earth • Groundtruth data	• Irrigated area

Table S2: Example EO-based methods used by researchers to assess agricultural intensity.

Indicator	Example Methods
Crop yield	<p>NB: A brief review of crop yield estimation techniques can be found in the introduction to Doraiswamy et al. (2005, 2004, 2003) and Kasampalis et al. (2018) provide an overview of crop growth models.</p> <ul style="list-style-type: none"> • Empirical regression-based modelling linking satellite-derived data (e.g. NDVI) to detailed official crop statistics (Becker-Reshef et al., 2010; Lobell et al., 2013; Salazar et al., 2007) or ground survey data (Ferencz et al., 2004; Ren et al., 2008; Yang et al., 2006); supplementary data may be included as additional explanatory variables (Balaghi et al., 2008; Prasad et al., 2006) • Estimate yield using regression models based on seasonal growth profiles from satellite-derived VIs (Kalubarme et al., 2003; Lai et al., 2018; Nagy et al., 2018; Son et al., 2014) • Crop yield simulation models incorporating satellite-derived data as either direct inputs or for calibration (Doraiswamy et al., 2005; Lobell et al., 2013; Moriondo et al., 2007) • Derive crop biomass using the Monteith light use efficiency approach (Awad, 2019; Leblon et al., 1991; Liu et al., 2010; Monteith, 1972; Morel et al., 2014; Pan et al., 2009; Patel et al., 2006)
Cropping area	<p>NB: Gallego (2004) provides a review of some common EO-based land cover area estimation techniques.</p> <ul style="list-style-type: none"> • Pixel counting & sub-pixel analysis (spectral unmixing, linear mixing models, mixture modelling) applied to classified satellite images; ground data used as an auxiliary/validation tool (Gallego, 2004; Gallego et al., 2014; Gumma et al., 2014; Vibhute and Gawali, 2013) • Regression analysis combining satellite-derived information with an accurate sample (e.g. ground survey data) (Gallego, 2004; Gallego et al., 2014; Vibhute and Gawali, 2013) • Derive a cropland probability layer using a combination of classified images and satellite-derived land surface phenology metrics (de Beurs and Ioffe, 2014) • SAR time-series decomposition (Canisius et al., 2018; Ponnurangam and Rao, 2018; Xu et al., 2019); SAR data may be integrated with multispectral data to assist crop classification (Gao et al., 2018; Shuai et al., 2019)
Cropping intensity – number of years a field is sown with crops and actually reaches harvest (de Beurs and Ioffe, 2014)	<p>Jain et al. (2013) provide a comparison of different methods to map cropping intensity. Methods included are:</p> <ul style="list-style-type: none"> • NDVI threshold method – define threshold for cropped land for a particular season based on satellite-derived NDVI, training data and regression tree analysis; use to classify pixels as cropped or uncropped agriculture for all seasons of interest • EVI peak method: (1) define threshold based on satellite-derived EVI and training data for cropped and non-cropped areas; (2) identify peaks in EVI time series; if peak exceeds threshold then classify it as cropped agriculture
Cropping frequency – number of years a pixel was cropped over an observation period (Estel et al., 2016)	<ul style="list-style-type: none"> • Hierarchical training technique (using EVI): (1) define the percent of each pixel cropped using higher resolution ground-truth imagery; (2) use this to calibrate EVI to quantify the percent of each pixel that was cropped in each season • Apply a series of simple decision rules to satellite-derived phenology metrics (phenology model) to distinguish cropped pixels from fallow lands. <i>Example rules in de Beurs & Ioffe (2014).</i>
Multi-cropping – number of harvests within a single year (i.e. growing season) (Estel et al., 2016)	<ul style="list-style-type: none"> • Use Temporal Mixture Analysis of end-member phenologies to determine whether a pixel is single, double or triple cropped (Jain et al., 2013) • Use a satellite-based phenology algorithm (e.g. Biradar and Xiao, 2011) to delineate the number of cropping cycles in a year • Apply time-series segmentation/iterative moving-window methodology to (smoothed) EVI time series to identify greening and browning phases and therefore cropping cycles; use to determine number of cropping cycles per year (Gray et al., 2014)
Cropping intensity – number of cropping cycles per year (L. Li et al., 2014)	<ul style="list-style-type: none"> • Use TIMESAT computer software to count the number of vegetation peaks in NDVI per growing season (Jönsson and Eklundh, 2004 in L. Li et al., 2014; Z. Li et al., 2014) • Determine the number of growth cycles in a year by counting the number of peaks (using thresholding techniques) on a crop growth curve based on satellite derived VI (e.g. EVI) (Yan et al., 2014)
Crop duration ratio	<p>Ratio of the time period (during the growing season) for which a pixel was cropped and the total length of the growing season (Estel et al., 2016)</p>

Table S2 Continued

Fallow cycles – recurring periods of fallow cropland	To identify fallow cycles: (1) Map active/fallow farmland based on NDVI time series (Estel <i>et al.</i> , 2015); (2) Filter time series for ‘chain segments’ i.e. certain number of consecutive fallow years; (3) count chain occurrence per pixel across entire time series; (4) summarise all chain segments using a weighting scheme (see Estel <i>et al.</i> , 2016 for details).
--	--

Table S3: Example EO-based methods used by researchers to assess vegetation health.

Indicator	Example EO-based Methods
Crop condition	<p><i>NB: A review of remote sensing methods for assessing crop condition can be found in Vibhute and Gawali (2013).</i></p> <ul style="list-style-type: none"> • Vegetation indices e.g. NDVI, NDWI, SAVI etc. – assume the higher the indices, the better the crop condition (Ali and Pelkey, 2013; Vibhute and Gawali, 2013) • Same-period comparing – compare EO-derived data (e.g. NDVI, LAI, VCI) of a specific period with data from period in history to determine areas of deterioration, no-change & improvement (Vibhute and Gawali, 2013; Wu <i>et al.</i>, 2015, 2014; Zhang <i>et al.</i>, 2014) • Crop growth profile monitoring (Jiang <i>et al.</i>, 2003; Vibhute and Gawali, 2013; Wu <i>et al.</i>, 2015, 2014; Zhang <i>et al.</i>, 2014)
Plant Trait Mapping	<i>NB: Homolova et al. (2013) and Andrew et al. (2014) provide reviews of remote sensing techniques for mapping various plant traits.</i>
Biophysical traits inc. biomass, fAPAR, photosynthetic capacity	<ul style="list-style-type: none"> • Empirical models (e.g. simple linear regression) relating limited field trait observations to EO-derived data such as vegetation indices (e.g. NDVI, EVI) & classified images (Baret and Guyot, 1991; Chen <i>et al.</i>, 2010; Homolová <i>et al.</i>, 2013; Jackson <i>et al.</i>, 2004; Karnieli <i>et al.</i>, 2013; Sakowska <i>et al.</i>, 2016; Schino <i>et al.</i>, 2003; Sibanda <i>et al.</i>, 2015; Turner <i>et al.</i>, 1999) • Radiative Transfer Models (RTM) (Homolová <i>et al.</i>, 2013; Myneni <i>et al.</i>, 1997) • Estimate fAPAR using Neural Networks (Baghdadi <i>et al.</i>, 2016) • Hyperspectral methods such as partial least squares regression (Hansen and Schjoerring, 2003)
Structural traits inc. crop/canopy height, leaf area index (LAI), biomass, canopy morphology	<p><i>NB: The introduction to Atzberger (2010) provides a brief overview of methods used to estimate LAI.</i></p> <ul style="list-style-type: none"> • Empirical models using spectral data, VIs or image texture metrics (Andrew <i>et al.</i>, 2014; Baret and Guyot, 1991; Clevers <i>et al.</i>, 2017; Delegido <i>et al.</i>, 2011; Z. Li <i>et al.</i>, 2014; Wulder <i>et al.</i>, 2004) • RTM-based approaches e.g. SAIL (Atzberger, 2010; Doraiswamy <i>et al.</i>, 2004; Frampton <i>et al.</i>, 2013; Homolová <i>et al.</i>, 2013; Jackson <i>et al.</i>, 2004; Myneni <i>et al.</i>, 1997; Verhoef, 1984) • Estimate using correlation between surface properties and backscatter from active sensors (Andrew <i>et al.</i>, 2014; Z. Li <i>et al.</i>, 2014) e.g. LiDAR (Drake <i>et al.</i>, 2002; van Leeuwen and Nieuwenhuis, 2010) & Radar (Brisco and Brown, 1998; Kasischke <i>et al.</i>, 1997)
Biochemical traits inc. chlorophyll (Ch) & water content, nitrogen (N) & phosphorous (P) status	<ul style="list-style-type: none"> • Empirical methods using vegetation indices (VIs) (Andrew <i>et al.</i>, 2014; Khanna <i>et al.</i>, 2007; Sakowska <i>et al.</i>, 2016) for example: Double-peak Canopy Nitrogen Index (DCNI) (Chen <i>et al.</i>, 2010); Modified triangle vegetation index 2 (Bagheri <i>et al.</i>, 2013); NDVI (Cheng <i>et al.</i>, 2008); Narrowband Green NDVI (NGNDVI) (Bausch and Khosla, 2010); Normalised Difference Water Index (NDWI) (Gao, 1996); Short-wave Infrared Water Stress Index (SIWSI) (Briant <i>et al.</i>, 2010; Fensholt and Sandholt, 2003); Triangular Greenness Index (TGI) (Hunt <i>et al.</i>, 2013); MERIS Terrestrial Chlorophyll Index (MTCI) (Dash and Curran, 2007) • RTM inversion e.g. REGFLEC (Andrew <i>et al.</i>, 2014; Boegh <i>et al.</i>, 2013; Frampton <i>et al.</i>, 2013; Homolová <i>et al.</i>, 2013; Houborg and Boegh, 2008; Jackson <i>et al.</i>, 2004; Trombetti <i>et al.</i>, 2008) • Estimate chlorophyll content using the red edge position (REP) (Z. Li <i>et al.</i>, 2014)

Table S4: Example EO-based methods used by researchers to assess soil quality and soil erosion/protection.

Indicator	Example EO-based Methods
Soil Quality	<i>NB: Shoshany et al. (2013) provide a review of EO methods for monitoring agricultural soil degradation</i>
Soil carbon (C) & organic matter (OM)	<ul style="list-style-type: none"> • Empirical modelling (e.g. partial least squares regression, random forest) (Andrew et al., 2014; Castaldi et al., 2019, 2016; Gholizadeh et al., 2018; Stevens et al., 2010) • Quantify using particular absorption features in the VIS-NIR-SWIR region (Ben Dor et al., 1999 in Shoshany et al., 2013) or the degree of concavity of the reflectance spectrum in VIS wavelengths (Andrew et al., 2014; Palacios-Orueta and Ustin, 1998)
Crop residue/conservation tillage density	<ul style="list-style-type: none"> • Map organic residue (or non-photosynthetic vegetation) cover using spectral unmixing approaches (Andrew et al., 2014; Pacheco and McNairn, 2010) • Spectral indices designed for detecting crop residues include: Normalised Difference Tillage Index (NDTI) and Normalised Difference Senescent Vegetation Index (NDSVI) (Daughtry et al., 2006, 2005) • Map crop residue using a multiband reflectance algorithm e.g. Crop Residue Index Multiband (CRIM) (Biard and Baret, 1997)
Nitrogen (N) status/availability	<ul style="list-style-type: none"> • Assess based on two premises: (1) N mineralisation & subsequent availability to growing crop will be proportional to OM content i.e. darker soil implies high soil nitrate levels (Scharf et al., 2002); (2) N stress increases canopy reflectance over all visible wavelengths (Beatty et al., 2000 in Scharf et al., 2002) – indices combining VIS & NIR regions may maximise sensitivity to N stress (Eitel et al., 2011; Scharf et al., 2002; Tilling et al., 2007) • Assess nitrogen status using surface indicators of subsurface nutrient conditions using multispectral and hyperspectral techniques (Shoshany et al., 2013)
Soil salinity	<p><i>NB: A review of the potentials and constraints of remote sensing-based soil salinity mapping can be found in Metternicht & Zinck (2003).</i></p> <ul style="list-style-type: none"> • Machine learning and regression-based models e.g. Multilayer Perception Neural Networks, Artificial Neural Networks, Gaussian Processes, Partial Least Square Regression, Support Vector Regression and Random Forest (Farifteh et al., 2007; Hoa et al., 2019; Taghadosi et al., 2019) • Categorical mapping of regions of differing soil salinity (e.g. high, medium, low) using hyperspectral satellite data and image classification (e.g. minimum distance, maximum likelihood) and spectral unmixing techniques (Ghosh et al., 2012; Hamzeh et al., 2016) • Quantitative mapping of soil salinity using indices derived from hyperspectral data and regression techniques (e.g. partial least squares regression, linear regression) (Bai et al., 2018; Hamzeh et al., 2013; Kumar et al., 2015; Mashimbye et al., 2012; Qian et al., 2019; Weng et al., 2008) • Distinguish ‘normal’ soil from moderately or severely salt-affected soils using brightness approach (Koshal 2010 in Shoshany et al., 2013) • Use spectral indices including Salinity Index (SI), Normalised Differential Salinity Index (NDSI) & Brightness Index (BI) (Asfaw et al., 2018; Dehni and Lounis, 2012; Khan et al., 2005; Shoshany et al., 2013) • Detect salinisation-related surface roughness features (e.g. crusting) using variation in radar backscattering & InSAR coherence signals (C, P & R wavelengths) (Metternicht and Zinck, 2003; Shoshany et al., 2013; Taylor et al., 1996)
Soil moisture (SM) content	<p><i>NB: Srivastava (2017) provides a review of satellite-based methods for monitoring soil moisture, while Petropoulos et al. (2018) provide an overview of the state of the art of EO techniques to derive operational estimates of soil moisture.</i></p> <ul style="list-style-type: none"> • Empirical & semi-empirical models relating backscattering coefficient to soil water content (& soil surface roughness) (Attarzadeh et al., 2018; Bao et al., 2018; Bousbih et al., 2018; Dubois et al., 1995; Genis et al., 2013; Hajj et al., 2017; Hosseini et al., 2015; Huang et al., 2019; Zhang et al., 2017) • RTM-based approaches relating soil dielectric constant to soil moisture (Babiet et al., 2018; Dubois et al., 1995; Hosseini et al., 2015; Jackson, 2002; Wagner et al., 2007) • Spectral indices e.g. Normalised Multiband Drought Index (NMDI) (Shoshany et al., 2013; Wang and Qu, 2009) • Retrieval from thermal data using apparent thermal inertia (ATI) (Shoshany et al., 2013; Verstraeten et al., 2006; Wang and Qu, 2009) • Map surface roughness & SM in sparsely vegetated landscapes using a multi-angle (radar-based) approach & an Integral Equation Model retrieval algorithm (Rahman et al., 2008)
Soil erosion/protection	<i>NB: Vrielling (2006) provide a review of satellite-based techniques for assessing erosion</i>
Vegetation cover	<ul style="list-style-type: none"> • Assess degree of protection based on amount of vegetation cover e.g. percentage ground cover, LAI (Cyr et al., 1995; Dwivedi et al., 1997; Fadul et al., 1999; Metternicht and Zinck, 1998; Wang et al., 2013)

Table S4 continued

Erosion feature detection	<ul style="list-style-type: none"> • Visual interpretation of high resolution images (Dwivedi <i>et al.</i>, 1997; Fadul <i>et al.</i>, 1999; Wang <i>et al.</i>, 2013) • Estimate metric dimensions & volume of individual patches of sheet, rill & gully erosion & densities (Metternicht and Zinck, 1998; Shoshany <i>et al.</i>, 2013) • Potential to use InSAR multi-temporal interferometric coherence change technique (Shoshany <i>et al.</i>, 2013)
Erosion modelling	<ul style="list-style-type: none"> • Erosion Potential Index (EPI) (Shoshany <i>et al.</i>, 2013) • Integrate EO-derived data into soil loss/erosion models (Cyr <i>et al.</i>, 1995) such as USLE (Universal Soil Loss Equation), ANSWERS (Areal Non-Point Source Watershed Environment Response Simulation), SEMMED (Soil Erosion Model for Mediterranean Regions) (De Jong, 1994; De Jong <i>et al.</i>, 1999; Ganasri and Ramesh, 2016; Shoshany <i>et al.</i>, 2013). EO derived-data includes: land use/land cover map (Baban and Yusof, 2001; De Jong <i>et al.</i>, 1999; Ganasri and Ramesh, 2016; Sharma and Singh, 1995), interception or total vegetation cover (De Jong, 1994), and soil parameter data (Baban and Yusof, 2001; De Jong <i>et al.</i>, 1999; Sharma and Singh, 1995)

Table S5: Example EO-based methods used by researchers to assess water quality and water availability.

Indicator	Example EO-based Methods
Water Quality	<i>NB: Reviews of various EO-based techniques for assessing various water quality parameters can be found in Gholizadeh et al. (2016a, 2016b); Dornhofer & Oppelt (2016) and Chang et al. (2015)</i>
Water Quality Indices	<ul style="list-style-type: none"> • Water Quality indices derived from different combinations of spectral bands (Vignolo <i>et al.</i>, 2006; Wen and Yang, 2010)
Physical water quality parameters: total suspended solids (TSS), turbidity, suspended sediment concentration (SSC), chlorophyll concentration, temperature & water clarity	<ul style="list-style-type: none"> • Empirical (simple or multiple regression) modelling – relate field data (or water quality indices derived from field data) to satellite data (e.g. band ratios) to estimate water quality parameters (Blix <i>et al.</i>, 2018; Carpenter and Carpenter, 1983; Chen <i>et al.</i>, 2007; Ha <i>et al.</i>, 2017; Hu <i>et al.</i>, 2004; Kloiber <i>et al.</i>, 2002; Lavery <i>et al.</i>, 1993; Liu <i>et al.</i>, 2017; Pereira-Sandoval <i>et al.</i>, 2018; Ritchie and Cooper, 2001; Sòria-perpinyà <i>et al.</i>, 2019) • Spectral unmixing-based approach – end-member spectra related to physical water quality parameters such as SSC (Martinez <i>et al.</i>, 2008)
Chemical water quality parameters: concentration of total nitrogen (TN), NO₃-N (nitrate as nitrogen) & total phosphorous (TP)	<ul style="list-style-type: none"> • Empirical (regression) modelling – relate field data to satellite data to estimate water quality parameters (Chen and Quan, 2012; Wu <i>et al.</i>, 2010) • Use neural network modelling (e.g. back-propagation neural network model) to establish a retrieval model for concentrations of TN & TP on the basis of satellite data (Xiao <i>et al.</i>, 2015)
Water quality proxy	<ul style="list-style-type: none"> • Assess health of vegetation alongside water bodies as a proxy for water quality, using vegetation indices (e.g. NDVI, EVI) (Trivero <i>et al.</i>, 2013) • Identification and mapping of submergent aquatic vegetation using image interpretation and classification techniques (Ackleson and Klemas, 1987; Dogan <i>et al.</i>, 2009; Wolter <i>et al.</i>, 2005; Yang, 2005)
Water Availability	
Water body area & configuration	<ul style="list-style-type: none"> • Detect/classify water bodies using optical data (NIR & SWIR regions) or spectral indices (e.g. NDVI & NDWI) (Andrew <i>et al.</i>, 2014; Frazier and Page, 2000; Mueller <i>et al.</i>, 2016; Smith, 1997; Tulbure and Broich, 2013) • WiPE water body classification algorithm (Ngoc <i>et al.</i>, 2019) • Determine water body area using pixel counting & vector-based GIS methods (Verpoorter <i>et al.</i>, 2012) • Quantify the spatial configuration of water bodies (e.g. number of water bodies, mean surface water body area) based on classified satellite images using FRAGSTATS software (v4) (McGarigal <i>et al.</i>, 2012; Tulbure and Broich, 2013) <p><i>NB: Detection of water bodies may be enhanced through use of techniques such as Principal Component Analysis (PCA) (Verpoorter et al., 2012)</i></p>
Water level & volume	<ul style="list-style-type: none"> • Estimate water level using satellite altimetry (Guo <i>et al.</i>, 2009; Koblinsky <i>et al.</i>, 1993; Michailovsky <i>et al.</i>, 2012; Smith, 1997; Sulistioadi <i>et al.</i>, 2015) • Use satellite-derived LC data as an input into models to estimate the volume of water yield available for consumptive purposes (Crossman <i>et al.</i>, 2013)

Table S5 continued

Water use efficiency & crop water stress	<ul style="list-style-type: none"> • Use satellite-derived data including VIs , NIR & TIR data as inputs into evapotranspiration models such as SEBAL (Surface Energy Balance for Land) (Bastiaanssen, 2000; Bastiaanssen <i>et al.</i>, 1998; Mutiga <i>et al.</i>, 2010), METRIC (Allen <i>et al.</i>, 2007) and ALEXI/DisALEXI (Anderson <i>et al.</i>, 2011) to predict actual evapotranspiration as an indicator of crop water stress and whether water is being used as intended (Gonzalez-Dugo <i>et al.</i>, 2009; Mutiga <i>et al.</i>, 2010) • Use TIR satellite data to calculate Evaporative Stress Index (ESI) to detect drought conditions and to infer crop health (Anderson and Kustas, 2008)
--	---

Table S6: Example EO-based methods used by researchers to assess biodiversity.

Indicator	Example EO-based Methods
<p><i>NB: EO-based techniques for monitoring biodiversity have been review by a number of authors including Gillespie et al. (2008), Kerr and Ostrovsky (2003), Kuenzer et al. (2014), Mairota et al. (2015), Nagendra (2001), Rocchini et al. (2010), Turner et al. (2003) and Wang et al. (2010)</i></p>	
Direct mapping of individuals and associations	<ul style="list-style-type: none"> • Map individual plants or associations of single species by applying pixel- or object-based classification procedures to high spatial resolution data (Ban, 2003; Clark <i>et al.</i>, 2001; Crossman <i>et al.</i>, 2013; Feng <i>et al.</i>, 2010; Gillespie <i>et al.</i>, 2008; Lauver, 1997; Nagendra and Gadgil, 1999; Turner <i>et al.</i>, 2003; Vibhute and Gawali, 2013). Species differentiation aided by differences in size, shape and vertical structure of canopies in active RS (e.g. LiDAR) or hyperspectral data (Andrew <i>et al.</i>, 2014) • Extract unique multi-temporal signature for different crops from VIs (e.g. NDVI & EVI) (Wardlow <i>et al.</i>, 2007) • Harmonic (Fourier) analysis of NDVI time series (Jakubauskas <i>et al.</i>, 2002; Mingwei <i>et al.</i>, 2008) • Classify vegetation types using visual and digital interpretation of false colour composites and SAR images based on derived characteristics including size, shape and texture (Blaes <i>et al.</i>, 2005; Ravan <i>et al.</i>, 1995)
Plant (and animal) species diversity	<ul style="list-style-type: none"> • Assess species diversity and distribution patterns by examining direct relationships between EO-derived spectral radiance values and species distribution patterns recorded from field observations (Feng <i>et al.</i>, 2010; Nagendra, 2001) • Use satellite-based land use and landscape complexity indices (e.g. patch shape indices) to predict regional plant species diversity (Honnay <i>et al.</i>, 2003) • Predict distribution or probability of occurrence of individual species and species assemblages using multiple regression analysis and EO-based data such as land cover maps (Jennings, 2000; Kerr <i>et al.</i>, 2001; Kerr and Ostrovsky, 2003; Luoto <i>et al.</i>, 2002a; Saveraid <i>et al.</i>, 2001); supplementary material such as climate and topography data may be incorporated (Cumming, 2000; Nagendra, 2001)
Habitat suitability	<ul style="list-style-type: none"> • Model distribution and abundance of single species using detailed information about known habitat requirements and EO-derived land cover, habitat maps and landscape metrics (Amici <i>et al.</i>, 2010; Feng <i>et al.</i>, 2010; Kerr <i>et al.</i>, 2001; Kerr and Ostrovsky, 2003; Z. Li <i>et al.</i>, 2014; Luoto <i>et al.</i>, 2002b; Mairota <i>et al.</i>, 2015; Nagendra, 2001; Weiers <i>et al.</i>, 2004) • Habitat suitability parameters include: spectral and textural indexes (Muñoz and Felicísimo, 2004; Stickler and Southworth, 2008); canopy cover (Davis <i>et al.</i>, 2007); NPP (Meynard and Quinn, 2007); existence of suitable water bodies (Weiers <i>et al.</i>, 2004); and hedgerow networks (Vannier <i>et al.</i>, 2011)
Landscape structure	<ul style="list-style-type: none"> • Derive quantitative measures of landscape structure (e.g. composition, isolation and complexity) from land cover classifications (Gustafson, 1998; Kuenzer <i>et al.</i>, 2014; Luoto <i>et al.</i>, 2002a; Rocchini <i>et al.</i>, 2010) • Landscape metrics can be computed by software products including FRAGSTATS (v4) using raster or vector data (McGarigal <i>et al.</i>, 2012) • Landscape diversity may be represented using diversity indices combining <i>richness</i> (number of classes present) and <i>evenness</i> (distribution of area among classes) (Gustafson, 1998). Examples include Shannon's and Simpson's diversity indices (Shannon and Weaver, 1948 and Simpson, 1949 in Gustafson, 1998). • Use image classification to map landscape connectivity elements (e.g. hedgerows) (Vannier <i>et al.</i>, 2011) • Quantify landscape fragmentation using pattern indices (Saura, 2004) such as number of patches and mean patch size (Turner and Ruscher, 1988 in Saura, 2004) and patch cohesion index (Schumaker, 1996)

Table S6 continued

<i>Species richness</i>	<ul style="list-style-type: none"> • Species-Energy Theory – species richness is proportional to NPP, derived from e.g. NDVI (Currie, 1991; Kerr and Ostrovsky, 2003; Z. Li <i>et al.</i>, 2014; Nagendra, 2001) • Spectral Variation Hypothesis – assume higher variation in spectra implies higher habitat heterogeneity, allowing coexistence of more species and consequently higher species richness (Diamond, 1988 in Fairbanks and McGwire, 2004; Z. Li <i>et al.</i>, 2014; Palmer <i>et al.</i>, 2002; Rocchini <i>et al.</i>, 2007) • Estimate spatial variation in species richness based on NDVI variability, vegetation classification map & multiple regression analysis (Bawa <i>et al.</i>, 2002; Bino <i>et al.</i>, 2008; Bonthoux <i>et al.</i>, 2018; Carrasco <i>et al.</i>, 2018; Fairbanks and McGwire, 2004; Gould, 2000)
<i>Invasive species</i>	<p>NB: Bradley <i>et al.</i> (2014) provide a review of remote sensing-based techniques for detecting invasive species</p> <ul style="list-style-type: none"> • Identify invasive species using visual interpretation, pixel-based & object-based classification, & spectral mixing/unmixing approaches (Huang and Asner, 2009; Z. Li <i>et al.</i>, 2014; Walsh <i>et al.</i>, 2008) • Map vegetation species from spectral and textural data using image classification techniques (e.g. maximum likelihood classification) (Kimothi and Dasari, 2010; Laba <i>et al.</i>, 2010, 2008; Mirik and Ansley, 2012), neural networks (Fuller, 2005) and principal component analysis (Tsai <i>et al.</i>, 2007) • Texture-augmented image analysis (Tsai and Chou, 2006) • Maximum Entropy Model (Evangelista <i>et al.</i>, 2009)

Table S7: Example EO-based methods used by researchers to assess ecosystem health.

Indicator	Example EO-based Methods	
	NB: EO-based techniques for monitoring ecosystem health have been reviewed by a number of authors including Andrew <i>et al.</i> (2014), Feng <i>et al.</i> (2010) and Z. Li <i>et al.</i> (2014)	
<i>Vigour</i>	<p><i>Net Primary Productivity (NPP) & Gross Primary Productivity (GPP)</i></p> <ul style="list-style-type: none"> • Model based on Light Use Efficiency (LUE) Concept (Feng <i>et al.</i>, 2010; Monteith, 1972 and Prince 1991 in Z. Li <i>et al.</i>, 2014; Ruimy and Saugier, 1994) • Statistical empirical model of GPP or NPP & a vegetation indices such as NDVI or EVI (Feng <i>et al.</i>, 2010; Z. Li <i>et al.</i>, 2014; Olofsson <i>et al.</i>, 2008; Xu <i>et al.</i>, 2012) • Estimate GPP based on photosynthetic capacity quantified using satellite-based leaf Ch content estimates e.g. from CI_{green} index (Gitelson <i>et al.</i>, 2008; Houborg <i>et al.</i>, 2013) 	
	<p><i>Fractional cover of green vegetation, non-photosynthetic vegetation (NPV) & bare soil</i></p> <ul style="list-style-type: none"> • Spectral mixing approach (Asner and Heidebrecht, 2002; Gill and Phinn, 2009; Gitelson, 2013; Z. Li <i>et al.</i>, 2014; Pacheco and McNairn, 2010) • Empirical model of fractional vegetation cover & vegetation indices (Carlson and Ripley, 1997; Gitelson, 2013; Guerschman <i>et al.</i>, 2009; Z. Li <i>et al.</i>, 2014; Wang <i>et al.</i>, 2018) • Estimate using a neural network based on NIR & red reflectances (Baret <i>et al.</i>, 1995; Gitelson, 2013) 	
	<p><i>Biochemical properties inc. N, P & chlorophyll</i></p> <ul style="list-style-type: none"> • Empirical modelling (based on biochemical spectra features) inc. simple linear regression, partial least-squares regression (PLSR) & stepwise linear regression (SMLR) (Homolová <i>et al.</i>, 2013; Z. Li <i>et al.</i>, 2014) 	
<i>Organisation</i>	<p><i>Species richness & biodiversity</i></p> <p>See table S6</p>	
	<p><i>Structural traits</i></p> <p>See table S2</p>	
<i>Resilience</i>	<ul style="list-style-type: none"> • Assessed based on a ratio of a given ES health indicator, e.g. aboveground biomass, measured pre- & post-disturbance (Z. Li <i>et al.</i>, 2014) • Vegetation indices e.g. NDVI time series frequently used to assess/monitor variation in vegetation health & deviation from normal conditions over time or in response to specific disturbances (Z. Li <i>et al.</i>, 2014) such as climate change (Li and Guo, 2012), wildfires (Díaz-Delgado <i>et al.</i>, 2002) & grazing intensity (Numata <i>et al.</i>, 2007) 	

Table S7 continued

Ecosystem Services as a Proxy for Ecosystem Health

- Use EO data (e.g. land cover) as an input for Ecosystem Services Models e.g. InVEST, ARIES, SolVES, GUMBO to assess the ability of an ecosystem to provide various ESS
- Indirect modelling techniques include deriving empirical models of ESS or their providers based on spatial environmental covariates, and using maps of biophysical drivers of ESS supply to parameterise mechanistic models (Andrew *et al.*, 2014).

References for Tables S1-S7

- Ackleson, S.G., Klemas, V., 1987. Remote sensing of submerged aquatic vegetation in lower chesapeake bay: A comparison of Landsat MSS to TM imagery. *Remote Sens. Environ.* 22, 235–248. doi:10.1016/0034-4257(87)90060-5
- Ali, R., Pelkey, N., 2013. Satellite images indicate vegetation degradation due to invasive herbivores in the Andaman Islands. *Curr. Sci.* 105, 209–214.
- Allen, R.G., Tasumi, M., Morse, A., Trezza, R., Wright, J.L., Bastiaanssen, W., Kramber, W., Lorite, I., Robison, C.W., 2007. Satellite-Based Energy Balance for Mapping Evapotranspiration with Internalized Calibration (METRIC) —Applications. *J. Irrig. Drain. Eng.* 133, 395–406. doi:10.1061/(ASCE)0733-9437(2007)133:4(395)
- Amici, V., Geri, F., Battisti, C., 2010. An integrated method to create habitat suitability models for fragmented landscapes. *J. Nat. Conserv.* 18, 215–223. doi:10.1016/j.jnc.2009.10.002
- Anderson, M., Kustas, W., 2008. Thermal remote sensing of drought and evapotranspiration. *Eos (Washington, DC)*. 89, 233–240. doi:10.1029/2008EO260001
- Anderson, M.C., Kustas, W.P., Norman, J.M., Hain, C.R., Mecikalski, J.R., Schultz, L., González-Dugo, M.P., Cammalleri, C., D’Urso, G., Pimstein, A., Gao, F., 2011. Mapping daily evapotranspiration at field to continental scales using geostationary and polar orbiting satellite imagery. *Hydrol. Earth Syst. Sci.* 15, 223–239. doi:10.5194/hess-15-223-2011
- Andrew, M.E., Wulder, M.A., Nelson, T.A., 2014. Potential contributions of remote sensing to ecosystem service assessments. *Prog. Phys. Geogr.* 38, 1–26. doi:10.1177/0309133314528942
- Armengot, L., José-María, L., Blanco-Moreno, J.M., Bassa, M., Chamorro, L., Xavier Sans, F., 2011. A novel index of land use intensity for organic and conventional farming of Mediterranean cereal fields. *Agron. Sustain. Dev.* 31, 699–707. doi:10.1007/s13593-011-0042-0
- Asfaw, E., Suryabagavan, K. V., Argaw, M., 2018. Soil salinity modeling and mapping using remote sensing and GIS: The case of Wonji sugar cane irrigation farm, Ethiopia. *J. Saudi Soc. Agric. Sci.* 17, 250–258. doi:10.1016/j.jssas.2016.05.003
- Asner, G.P., Heidebrecht, K.B., 2002. Spectral unmixing of vegetation, soil and dry carbon cover in arid regions: comparing multispectral and hyperspectral observations. *Int. J. Remote Sens.* 23, 3939–3958. doi:10.1080/01431160110115960
- Attarzadeh, R., Amini, J., Notarnicola, C., Greifeneder, F., 2018. Synergetic Use of Sentinel-1 and Sentinel-2 Data for Soil Moisture Mapping at Plot Scale. *Remote Sens.* 10, 1285. doi:10.3390/rs10081285
- Atzberger, C., 2010. Inverting the PROSAIL canopy reflectance model using neural nets trained on streamlined databases. *J. Spectr. Imaging* 1, 1–13. doi:10.1255/jsi.2010.a2
- Awad, M., 2019. Toward Precision in Crop Yield Estimation Using Remote Sensing and Optimization Techniques. *Agriculture* 9, 54. doi:10.3390/agriculture9030054
- Baban, S.M.J., Yusof, K.W., 2001. Modelling soil erosion in tropical environments using remote sensing and geographical information systems. *Hydrol. Sci. -Journal-des Sci. Hydrol.* 46, 191–198. doi:10.1080/02626660109492815
- Bablt, A., Vu, P.V.H., Jacquemoud, S., Viallefont-Robinet, F., Fabre, S., Briottet, X., Sadeghi, M., Whiting, M.L., Baret, F., Tian, J., 2018. MARMIT: A multilayer radiative transfer model of soil reflectance to estimate surface soil moisture content in the solar domain (400–2500 nm). *Remote Sens. Environ.* 217, 1–17. doi:10.1016/j.rse.2018.07.031
- Baghdadi, N., Hajj, M., El, Zribi, M., Fayad, I., 2016. Coupling SAR C-band and optical data for soil moisture and leaf area index retrieval over irrigated grasslands. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* 9, 1229–1243.

- Bagheri, N., Ahmadi, H., Kazem Alavipanah, S., Omid, M., 2013. Multispectral remote sensing for site-specific nitrogen fertilizer management. *Pesqui. Agropeuária Bras.* 48, 1394–1401. doi:10.1590/S0100-204X2013001000011
- Bai, L., Wang, C., Zang, S., Wu, C., Luo, J., Wu, Y., 2018. Mapping soil alkalinity and salinity in northern songnen plain, China with the hj-1 hyperspectral imager data and partial least squares regression. *Sensors* 18, 3855. doi:10.3390/s18113855
- Balaghi, R., Tychon, B., Eerens, H., Jlibene, M., 2008. Empirical regression models using NDVI, rainfall and temperature data for the early prediction of wheat grain yields in Morocco. *Int. J. Appl. Earth Obs. Geoinf.* 10, 438–452. doi:10.1016/j.jag.2006.12.001
- Ban, Y., 2003. Synergy of multitemporal ERS-1 SAR and Landsat TM data for classification of agricultural crops. *Can. J. Remote Sens.* 29, 518–526. doi:10.5589/m03-014
- Bao, Y., Lin, L., Wu, S., Kwal Deng, K.A., Petropoulos, G.P., 2018. Surface soil moisture retrievals over partially vegetated areas from the synergy of Sentinel-1 and Landsat 8 data using a modified water-cloud model. *Int. J. Appl. Earth Obs. Geoinf.* 72, 76–85. doi:10.1016/j.jag.2018.05.026
- Baret, F., Clevers, J.G.P.W., Steven, M.D., 1995. The robustness of canopy gap fraction estimates from red and near-infrared reflectances: A comparison of approaches. *Remote Sens. Environ.* 54, 141–151. doi:10.1016/0034-4257(95)00136-O
- Baret, F., Guyot, G., 1991. Potentials and Limits of Vegetation Indices for LAI and APAR Assessment. *Remote Sens. Environ.* 35, 161–173. doi:10.1016/0034-4257(91)90009-U
- Bastiaanssen, W.G.M., 2000. SEBAL-based sensible and latent heat fluxes in the irrigated Gediz Basin, Turkey. *J. Hydrol.* 229, 87–100. doi:10.1016/S0022-1694(99)00202-4
- Bastiaanssen, W.G.M., Menenti, M., Feddes, R.A., Holtslag, A.A.M., 1998. A remote sensing surface energy balance algorithm for land (SEBAL) 1. Formulation. *J. Hydrol.* 212–213, 198–212. doi:10.1016/S0022-1694(98)00253-4
- Bausch, W.C., Khosla, R., 2010. QuickBird satellite versus ground-based multi-spectral data for estimating nitrogen status of irrigated maize. *Precis. Agric.* 11, 274–290. doi:10.1007/s11119-009-9133-1
- Bawa, K., Rose, J., Ganeshaiah, K.N., Barve, N., Kiran, M.C., Umashaanker, R., 2002. Assessing biodiversity from space: An example from the Western Ghats, India. *Conserv. Ecol.* 6, 7. doi:10.5751/ES-00434-060207
- Becker-Reshef, I., Vermote, E., Lindeman, M., Justice, C., 2010. A generalized regression-based model for forecasting winter wheat yields in Kansas and Ukraine using MODIS data. *Remote Sens. Environ.* 114, 1312–1323. doi:10.1016/j.rse.2010.01.010
- Biard, F., Baret, F., 1997. Crop Residue Estimation Using Multiband Reflectance. *Remote Sens. Environ.* 59, 530–536. doi:10.1016/S0034-4257(96)00125-3
- Bino, G., Levin, N., Darawshi, S., Van Der Hal, N., Reich-Solomon, a., Kark, S., 2008. Accurate prediction of bird species richness patterns in an urban environment using Landsat-derived NDVI and spectral unmixing. *Int. J. Remote Sens.* 29, 3675–3700. doi:10.1080/01431160701772534
- Biradar, C.M., Xiao, X., 2011. Quantifying the area and spatial distribution of double-and triple-cropping croplands in India with multi-temporal MODIS imagery in 2005. *Int. J. Remote Sens.* 32, 367–386. doi:10.1080/01431160903464179
- Blaes, X., Vanhalle, L., Defourny, P., 2005. Efficiency of crop identification based on optical and SAR image time series. *Remote Sens. Environ.* 96, 352–365. doi:10.1016/j.rse.2005.03.010
- Blix, K., Pálffy, K., Tóth, V.R., Eltoft, T., 2018. Remote sensing of water quality parameters over Lake Balaton by using Sentinel-3 OLCI. *Water* 10, 1428. doi:10.3390/w10101428
- Boegh, E., Houborg, R., Bienkowski, J., Braban, C.F., Dalgaard, T., Van Dijk, N., Dragosits, U., Holmes, E., Magliulo, V., Schelde, K., Di Tommasi, P., Vitale, L., Theobald, M.R., Cellier, P., Sutton, M.A., 2013. Remote sensing of LAI, chlorophyll and leaf nitrogen pools of crop- and grasslands in five European landscapes. *Biogeosciences* 10, 6279–6307. doi:10.5194/bg-10-6279-2013
- Bonthoux, S., Lefèvre, S., Herrault, P.-A., Sheeren, D., 2018. Spatial and Temporal Dependency of NDVI Satellite Imagery in Predicting Bird Diversity over France. *Remote Sens.* 10, 1136. doi:10.3390/rs10071136

- Bousbih, S., Zribi, M., Hajj, M. El, Baghdadi, N., Lili-Chabaane, Z., Gao, Q., Fanise, P., 2018. Soil moisture and irrigation mapping in a semi-arid region, based on the synergetic use of Sentinel-1 and Sentinel-2 data. *Remote Sens.* 10, 1953. doi:10.3390/rs10121953
- Bradley, B.A., 2014. Remote detection of invasive plants: A review of spectral, textural and phenological approaches. *Biol. Invasions* 16, 1411–1425. doi:10.1007/s10530-013-0578-9
- Briant, G., Gond, V., Laurance, S.G.W., 2010. Habitat fragmentation and the desiccation of forest canopies: A case study from eastern Amazonia. *Biol. Conserv.* 143, 2763–2769. doi:10.1016/j.biocon.2010.07.024
- Brisco, B., Brown, R.J., 1998. Agricultural Applications with Radar, in: Henderson, F.M., Lewis, A.J. (Eds.), *Principles and Applications of Imaging Radar - Manual of Remote Sensing Third Edition, Volume 2*. John Wiley & Sons, New York, pp. 381–406.
- Canisius, F., Shang, J., Liu, J., Huang, X., Ma, B., Jiao, X., Geng, X., Kovacs, J.M., Walters, D., 2018. Tracking crop phenological development using multi-temporal polarimetric Radarsat-2 data. *Remote Sens. Environ.* 210, 508–518. doi:10.1016/j.rse.2017.07.031
- Carlson, T.N., Ripley, D.A., 1997. On the relation between NDVI, fractional vegetation cover, and leaf area index. *Remote Sens. Environ.* 62, 241–252. doi:10.1016/S0034-4257(97)00104-1
- Carpenter, D.J., Carpenter, S.M., 1983. Modeling Inland Water Quality Using Landsat Data. *Remote Sens. Environ.* 13, 345–352. doi:10.1016/0034-4257(83)90035-4
- Carrasco, L., Norton, L., Henrys, P., Siriwardena, G.M., Rhodes, C.J., Rowland, C., Morton, D., 2018. Habitat diversity and structure regulate British bird richness: Implications of non-linear relationships for conservation. *Biol. Conserv.* 226, 256–263. doi:10.1016/j.biocon.2018.08.010
- Castaldi, F., Hueni, A., Chabrilat, S., Ward, K., Buttafuoco, G., Bomans, B., Vreys, K., Brell, M., van Wesemael, B., 2019. Evaluating the capability of the Sentinel 2 data for soil organic carbon prediction in croplands. *ISPRS J. Photogramm. Remote Sens.* 147, 267–282. doi:10.1016/j.isprsjprs.2018.11.026
- Castaldi, F., Palombo, A., Santini, F., Pascucci, S., Pignatti, S., Casa, R., 2016. Evaluation of the potential of the current and forthcoming multispectral and hyperspectral imagers to estimate soil texture and organic carbon. *Remote Sens. Environ.* 179, 54–65. doi:10.1016/j.rse.2016.03.025
- Chang, N. Bin, Imen, S., Vannah, B., 2015. Remote sensing for monitoring surface water quality status and ecosystem state in relation to the nutrient cycle: A 40-year perspective. *Crit. Rev. Environ. Sci. Technol.* 45, 101–166. doi:10.1080/10643389.2013.829981
- Chen, J., Quan, W., 2012. Using Landsat/TM imagery to estimate nitrogen and phosphorus concentration in Taihu Lake, China. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* 5, 273–280. doi:10.1109/JSTARS.2011.2174339
- Chen, P., Haboudane, D., Tremblay, N., Wang, J., Vigneault, P., Li, B., 2010. New spectral indicator assessing the efficiency of crop nitrogen treatment in corn and wheat. *Remote Sens. Environ.* 114, 1987–1997. doi:10.1016/j.rse.2010.04.006
- Chen, Z., Muller-Karger, F.E., Hu, C., 2007. Remote sensing of water clarity in Tampa Bay. *Remote Sens. Environ.* 109, 249–259. doi:10.1016/j.rse.2007.01.002
- Cheng, Y.B., Ustin, S.L., Riaño, D., Vanderbilt, V.C., 2008. Water content estimation from hyperspectral images and MODIS indexes in Southeastern Arizona. *Remote Sens. Environ.* 112, 363–374. doi:10.1016/j.rse.2007.01.023
- Clark, P.E., Seyfried, M.S., Harris, B., 2001. Intermountain plant community classification using Landsat TM and SPOT HRV data. *J. Range Manag.* 54, 152–160. doi:10.2307/4003176
- Clevers, J.G.P.W., Kooistra, L., van den Brande, M.M.M., 2017. Using Sentinel-2 data for retrieving LAI and leaf and canopy chlorophyll content of a potato crop. *Remote Sens.* 9, 405. doi:10.3390/rs9050405
- Crossman, N.D., Burkhard, B., Nedkov, S., Willems, L., Petz, K., Palomo, I., Drakou, E.G., Martín-Lopez, B., McPhearson, T., Boyanova, K., Alkemade, R., Egoh, B., Dunbar, M.B., Maes, J., 2013. A blueprint for mapping and modelling ecosystem services. *Ecosyst. Serv.* 4, 4–14. doi:10.1016/j.ecoser.2013.02.001
- Cumming, G.S., 2000. Using habitat models to map diversity: Pan-African species richness of ticks (Acari: Ixodida). *J. Biogeogr.* 27, 425–440. doi:10.1046/j.1365-2699.2000.00419.x

- Currie, D.J., 1991. Energy and Large-Scale Patterns of Animal-and Plant-Species Richness. *Am. Nat.* 137, 27–49. doi:10.1086/285144
- Cyr, L., Bonn, F., Pesant, A., 1995. Vegetation indices derived from remote sensing for an estimation of soil protection against water erosion. *Ecol. Modell.* 79, 277–285. doi:10.1016/0304-3800(94)00182-H
- Dash, J., Curran, P.J., 2007. Evaluation of the MERIS terrestrial chlorophyll index (MTCI). *Adv. Sp. Res.* 39, 100–104. doi:10.1016/j.asr.2006.02.034
- Daughtry, C.S.T., Doraiswamy, P.C., Hunt, E.R., Stern, A.J., McMurtrey Iii, J.E., Prueger, J.H., 2006. Remote sensing of crop residue cover and soil tillage intensity. *Soil Tillage Res.* 91, 101–108. doi:10.1016/j.still.2005.11.013
- Daughtry, C.S.T., Hunt, E.R., Doraiswamy, P.C., McMurtrey, J.E., 2005. Remote sensing the spatial distribution of crop residues. *Agron. J.* 97, 864–871. doi:10.2134/agronj2003.0291
- Davis, F.W., Seo, C., Zielinski, W.J., 2007. Regional variation in home-range-scale habitat models for fisher (*Martes pennanti*) in California. *Ecol. Appl.* 17, 2195–2213. doi:10.1890/06-1484.1
- de Beurs, K.M., Ioffe, G., 2014. Use of Landsat and MODIS data to remotely estimate Russia's sown area. *J. Land Use Sci.* 9, 377–401. doi:10.1080/1747423X.2013.798038
- De Jong, S.M., 1994. Derivation of vegetative variables from a landsat TM image for modelling soil erosion. *Earth Surf. Process. Landforms* 19, 165–178. doi:10.1002/esp.3290190207
- De Jong, S.M., Paracchini, M.L., Bertolo, F., Folving, S., Megier, J., De Roo, A.P.J., 1999. Regional assessment of soil erosion using the distributed model SEMMED and remotely sensed data. *Catena* 37, 291–308. doi:10.1016/S0341-8162(99)00038-7
- Dehni, A., Lounis, M., 2012. Remote sensing techniques for salt affected soil mapping: Application to the Oran region of Algeria. *Procedia Eng.* 33, 188–198. doi:10.1016/j.proeng.2012.01.1193
- Delegido, J., Verrelst, J., Alonso, L., Moreno, J., 2011. Evaluation of sentinel-2 red-edge bands for empirical estimation of green LAI and chlorophyll content. *Sensors* 11, 7063–7081. doi:10.3390/s110707063
- Díaz-Delgado, R., Lloret, F., Pons, X., Terradas, J., 2002. Satellite evidence of decreasing resilience in Mediterranean plant communities after recurrent wildfires. *Ecology* 83, 2293–2303. doi:10.1890/0012-9658(2002)083[2293:SEODRI]2.0.CO;2
- Dogan, O.K., Akyurek, Z., Beklioglu, M., 2009. Identification and mapping of submerged plants in a shallow lake using quickbird satellite data. *J. Environ. Manage.* 90, 2138–2143. doi:10.1016/j.jenvman.2007.06.022
- Donald, P.F., Green, R.E., Heath, M.F., 2001. Agricultural intensification and the collapse of Europe's farmland bird populations. *Proc. R. Soc. B* 268, 25–29. doi:10.1098/rspb.2000.1325
- Doraiswamy, P.C., Hatfield, J.L., Jackson, T.J., Akhmedov, B., Prueger, J., Stern, A., 2004. Crop condition and yield simulations using Landsat and MODIS. *Remote Sens. Environ.* 92, 548–559. doi:10.1016/j.rse.2004.05.017
- Doraiswamy, P.C., Moulin, S., Cook, P.W., Stern, A., 2003. Crop Yield Assessment from Remote Sensing. *Photogramm. Eng. Remote Sens.* 69, 665–674. doi:10.14358/pers.69.6.665
- Doraiswamy, P.C., Sinclair, T.R., Hollinger, S., Akhmedov, B., Stern, A., Prueger, J., 2005. Application of MODIS derived parameters for regional crop yield assessment. *Remote Sens. Environ.* 97, 192–202. doi:10.1016/j.rse.2005.03.015
- Dörnhöfer, K., Oppelt, N., 2016. Remote sensing for lake research and monitoring - Recent advances. *Ecol. Indic.* 64, 105–122. doi:10.1016/j.ecolind.2015.12.009
- Drake, J.B., Dubayah, R.O., Clark, D.B., Knox, R.G., Blair, J.B., Hofton, M.A., Chazdon, R.L., Weishampel, J.F., Prince, S., 2002. Estimation of tropical forest structural characteristics, using large-footprint lidar. *Remote Sens. Environ.* 79, 305–319. doi:10.1016/S0034-4257(01)00281-4
- Dubois, P.C., van Zyl, J., Engman, T., 1995. Measuring Soil Moisture with Imaging Radars. *IEEE Trans. Geosci. Remote Sens.* 33, 915–926. doi:10.1109/36.406677
- Dwivedi, R.S., Sankar, T.R., Venkataratnam, L., Karale, R.L., Gawande, S.P., Rao, K.V.S., Senchaudhary, S., Bhaumik, K.R., Mukharjee, K.K., 1997. The inventory and monitoring of eroded lands using remote sensing data. *Int. J. Remote Sens.* 18, 107–119. doi:10.1080/014311697219303

- Egorov, E., Prati, D., Durka, W., Michalski, S., Fischer, M., Schmitt, B., Blaser, S., Brändle, M., 2014. Does land-use intensification decrease plant phylogenetic diversity in local grasslands? *PLoS One* 9, e103252. doi:10.1371/journal.pone.0103252
- Eitel, J.U.H., Vierling, L.A., Litvak, M.E., Long, D.S., Schulthess, U., Ager, A.A., Krofcheck, D.J., Stoscheck, L., 2011. Broadband, red-edge information from satellites improves early stress detection in a New Mexico conifer woodland. *Remote Sens. Environ.* 115, 3640–3646. doi:10.1016/j.rse.2011.09.002
- Estel, S., Kuemmerle, T., Alcántara, C., Levers, C., Prishchepov, A., Hostert, P., 2015. Mapping farmland abandonment and recultivation across Europe using MODIS NDVI time series. *Remote Sens. Environ.* 163. doi:10.1016/j.rse.2015.03.028
- Estel, S., Kuemmerle, T., Levers, C., Baumann, M., Hostert, P., 2016. Mapping cropland-use intensity across Europe using MODIS NDVI time series. *Environ. Res. Lett.* 11. doi:10.1088/1748-9326/11/2/024015
- Evangelista, P.H., Stohlgren, T.J., Morisette, J.T., Kumar, S., 2009. Mapping invasive tamarisk (*Tamarix*): A comparison of single-scene and time-series analyses of remotely sensed data. *Remote Sens.* 1, 519–533. doi:10.3390/rs1030519
- Fadul, H.M., Salih, A.A., Ali, I.A., Inanaga, S., 1999. Use of remote sensing to map gully erosion along the Atbara River, Sudan. *JAG* 1, 314–1999. doi:10.1016/S0303-2434(99)85010-7
- Fairbanks, D.H.K., McGwire, K.C., 2004. Patterns of floristic richness in vegetation communities of California: Regional scale analysis with multi-temporal NDVI. *Glob. Ecol. Biogeogr.* 13, 221–235. doi:10.1111/j.1466-822X.2004.00092.x
- Farifteh, J., Van der Meer, F., Atzberger, C., Carranza, E.J.M., 2007. Quantitative analysis of salt-affected soil reflectance spectra: A comparison of two adaptive methods (PLSR and ANN). *Remote Sens. Environ.* 110, 59–78. doi:10.1016/j.rse.2007.02.005
- Feng, X., Fu, B., Yang, X., Lü, Y., 2010. Remote sensing of ecosystem services: An opportunity for spatially explicit assessment. *Chinese Geogr. Sci.* 20, 522–535. doi:10.1007/s11769-010-0428-y
- Fensholt, R., Sandholt, I., 2003. Derivation of a shortwave infrared water stress index from MODIS near- and shortwave infrared data in a semiarid environment. *Remote Sens. Environ.* 87, 111–121. doi:10.1016/j.rse.2003.07.002
- Ferencz, C., Bognár, P., Lichtenberger, J., Hamar, D., Tarcsai, G., Timár, G., Molnár, G., Pásztor, S.Z., Steinbach, P., Székely, B., Ferencz, O.E., Ferencz-Árkos, I., 2004. Crop yield estimation by satellite remote sensing. *Int. J. Remote Sens.* 25, 4113–4149. doi:10.1080/01431160410001698870
- Frampton, W.J., Dash, J., Watmough, G., Milton, E.J., 2013. Evaluating the capabilities of Sentinel-2 for quantitative estimation of biophysical variables in vegetation. *ISPRS J. Photogramm. Remote Sens.* 82, 83–92. doi:10.1016/j.isprsjprs.2013.04.007
- Frazier, P.S., Page, K.J., 2000. Water body detection and delineating with Landsat TM data. *Photogramm. Eng. Remote Sens.* 66, 1461–1467.
- Fuller, D.O., 2005. Remote detection of invasive *Melaleuca* trees (*Melaleuca quinquenervia*) in South Florida with multispectral IKONOS imagery. *Int. J. Remote Sens.* 26, 1057–1063. doi:10.1080/01430060512331314119
- Gallego, F.J., 2004. Remote sensing and land cover area estimation. *Int. J. Remote Sens.* 25, 3019–3047. doi:10.1080/01431160310001619607
- Gallego, F.J., Kussul, N., Skakun, S., Kravchenko, O., Shelestov, A., Kussul, O., 2014. Efficiency assessment of using satellite data for crop area estimation in Ukraine. *Int. J. Appl. Earth Obs. Geoinf.* 29, 22–30. doi:10.1016/j.jag.2013.12.013
- Ganasri, B.P., Ramesh, H., 2016. Assessment of soil erosion by RUSLE model using remote sensing and GIS - A case study of Nethravathi Basin. *Geosci. Front.* 7, 953–961. doi:10.1016/j.gsf.2015.10.007
- Gao, B.C., 1996. NDWI - A normalized difference water index for remote sensing of vegetation liquid water from space. *Remote Sens. Environ.* 58, 257–266. doi:10.1016/S0034-4257(96)00067-3
- Gao, H., Wang, C., Wang, G., Zhu, J., Tang, Y., Shen, P., Zhu, Z., 2018. A crop classification method integrating GF-3 PolSAR and sentinel-2A optical data in the Dongting lake basin. *Sensors* 18, 3139. doi:10.3390/s18093139
- Genis, A., Vulfson, L., Blumberg, D.G., Sprinstin, M., Kotlyar, A., Freilikher, V., Ben-Asher, J., 2013. Retrieving parameters of bare soil surface roughness and soil water content under arid environment from ERS-1, -2 SAR data. *Int. J. Remote Sens.* 34, 6202–6215. doi:10.1080/01431161.2013.793862
- Gholizadeh, A., Žižala, D., Saberioon, M., Borůvka, L., 2018. Soil organic carbon and texture retrieving and mapping using

- proximal, airborne and Sentinel-2 spectral imaging. *Remote Sens. Environ.* 218, 89–103. doi:10.1016/j.rse.2018.09.015
- Gholizadeh, M.H., Melesse, A.M., Reddi, L., 2016a. A comprehensive review on water quality parameters estimation using remote sensing techniques. *Sensors* 16, 1298. doi:10.3390/s16081298
- Gholizadeh, M.H., Melesse, A.M., Reddi, L., 2016b. Spaceborne and airborne sensors in water quality assessment. *Int. J. Remote Sens.* 37, 3143–3180. doi:10.1080/01431161.2016.1190477
- Ghosh, G., Kumar, S., Saha, S.K., 2012. Hyperspectral Satellite Data in Mapping Salt-Affected Soils Using Linear Spectral Unmixing Analysis. *J. Indian Soc. Remote Sens.* 40, 129–136. doi:10.1007/s12524-011-0143-x
- Gill, T.K., Phinn, S.R., 2009. Improvements to ASTER-Derived Fractional Estimates of Bare Ground in a Savanna Rangeland. *IEEE Trans. Geosci. Remote Sens.* 47, 662–670. doi:10.1109/TGRS.2008.2004628
- Gillespie, T.W., Foody, G.M., Rocchini, D., Giorgi, A.P., Saatchi, S., 2008. Measuring and modelling biodiversity from space. *Prog. Phys. Geogr.* 32, 203–221. doi:10.1177/0309133308093606
- Gitelson, A.A., 2013. Remote estimation of crop fractional vegetation cover: the use of noise equivalent as an indicator of performance of vegetation indices. *Int. J. Remote Sens.* 34, 6054–6066. doi:10.1080/01431161.2013.793868
- Gitelson, A.A., Viña, A., Masek, J.G., Verma, S.B., Suyker, A.E., 2008. Synoptic monitoring of gross primary productivity of maize using landsat data. *IEEE Geosci. Remote Sens. Lett.* 5, 133–137. doi:10.1109/LGRS.2008.915598
- Gonzalez-Dugo, M.P., Neale, C.M.U., Mateos, L., Kustas, W.P., Prueger, J.H., Anderson, M.C., Li, F., 2009. A comparison of operational remote sensing-based models for estimating crop evapotranspiration. *Agric. For. Meteorol.* 149, 1843–1853. doi:10.1016/j.agrformet.2009.06.012
- Gould, W., 2000. Remote Sensing of Vegetation, Plant Species Richness, and Regional Biodiversity Hotspots. *Ecol. Appl.* 10, 1861–1870. doi:10.2307/2641244
- Gray, J., Friedl, M., Frohling, S., Ramankutty, N., Nelson, A., Gumma, M.K., 2014. Mapping Asian cropping intensity with MODIS. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* 7, 3373–3379. doi:10.1109/JSTARS.2014.2344630
- Guerschman, J.P., Hill, M.J., Renzullo, L.J., Barrett, D.J., Marks, A.S., Botha, E.J., 2009. Estimating fractional cover of photosynthetic vegetation, non-photosynthetic vegetation and bare soil in the Australian tropical savanna region upscaling the EO-1 Hyperion and MODIS sensors. *Remote Sens. Environ.* 113, 928–945. doi:10.1016/j.rse.2009.01.006
- Gumma, M.K., Thenkabail, P.S., Maunahan, A., Islam, S., Nelson, A., 2014. Mapping seasonal rice cropland extent and area in the high cropping intensity environment of Bangladesh using MODIS 500m data for the year 2010. *ISPRS J. Photogramm. Remote Sens.* 91, 98–113. doi:10.1016/j.isprsjprs.2014.02.007
- Guo, J., Chang, X., Gao, Y., Sun, J., Hwang, C., 2009. Lake level variations monitored with satellite altimetry waveform retracking. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* 2, 80–86. doi:10.1109/JSTARS.2009.2021673
- Gustafson, E.J., 1998. Quantifying Landscape Spatial Pattern: What Is the State of the Art? *Ecosystems* 1, 143–156. doi:10.1007/s100219900011
- Ha, N.T.T., Thao, N.T.P., Koike, K., Nhuan, M.T., 2017. Selecting the Best Band Ratio to Estimate Chlorophyll-a Concentration in a Tropical Freshwater Lake Using Sentinel 2A Images from a Case Study of Lake Ba Be (Northern Vietnam). *ISPRS Int. J. Geo-Information* 6, 290. doi:10.3390/ijgi6090290
- Hajj, M. El, Baghdadi, N., Zribi, M., Bazzi, H., 2017. Synergic use of Sentinel-1 and Sentinel-2 images for operational soil moisture mapping at high spatial resolution over agricultural areas. *Remote Sens.* 9, 1292. doi:10.3390/rs9121292
- Hamzeh, S., Naseri, A.A., AlaviPanah, S.K., Bartholomeus, H., Herold, M., 2016. Assessing the accuracy of hyperspectral and multispectral satellite imagery for categorical and quantitative mapping of salinity stress in sugarcane fields. *Int. J. Appl. Earth Obs. Geoinf.* 52, 412–421. doi:10.1016/j.jag.2016.06.024
- Hamzeh, S., Naseri, A.A., AlaviPanah, S.K., Mojaradi, B., Bartholomeus, H.M., Clevers, J.G.P.W., Behzad, M., 2013. Estimating salinity stress in sugarcane fields with spaceborne hyperspectral vegetation indices. *Int. J. Appl. Earth Obs. Geoinf.* 21, 282–290. doi:10.1016/j.jag.2012.07.002
- Hansen, P.M., Schjoerring, J.K., 2003. Reflectance measurement of canopy biomass and nitrogen status in wheat crops using normalized difference vegetation indices and partial least squares regression. *Remote Sens. Environ.* 86, 542–553.

- Herzog, F., Steiner, B., Bailey, D., Baudry, J., Billeter, R., Bukáček, R., De Blust, G., De Cock, R., Dirksen, J., Dormann, C.F., De Filippi, R., Frossard, E., Liira, J., Schmidt, T., Stöckli, R., Thenail, C., Van Wingerden, W., Bugter, R., 2006. Assessing the intensity of temperate European agriculture at the landscape scale. *Eur. J. Agron.* 24, 165–181. doi:10.1016/j.eja.2005.07.006
- Hoa, P.V., Giang, N.V., Binh, N.A., Hai, L.V.H., Pham, T.D., Hasanlou, M., Bui, D.T., 2019. Soil salinity mapping using SAR Sentinel-1 data and advanced machine learning algorithms: A case study at Ben Tre Province of the Mekong River Delta (Vietnam). *Remote Sens.* 11, 128. doi:10.3390/rs11020128
- Homolová, L., Malenovský, Z., Clevers, J.G.P.W., García-Santos, G., Schaepman, M.E., 2013. Review of optical-based remote sensing for plant trait mapping. *Ecol. Complex.* 15, 1–16. doi:10.1016/j.ecocom.2013.06.003
- Honnay, O., Piessens, K., Van Landuyt, W., Hermy, M., Gulinck, H., 2003. Satellite based land use and landscape complexity indices as predictors for regional plant species diversity. *Landsc. Urban Plan.* 63, 241–250. doi:10.1016/S0169-2046(02)00194-9
- Hosseini, R., Newlands, N.K., Dean, C.B., Takemura, A., 2015. Statistical modeling of soil moisture, integrating satellite remote-sensing (SAR) and ground-based data. *Remote Sens.* 7, 2752–2780. doi:10.3390/rs70302752
- Houborg, R., Boegh, E., 2008. Mapping leaf chlorophyll and leaf area index using inverse and forward canopy reflectance modeling and SPOT reflectance data. *Remote Sens. Environ.* 112, 186–202. doi:10.1016/j.rse.2007.04.012
- Houborg, R., Cescatti, A., Migliavacca, M., Kustas, W.P., 2013. Satellite retrievals of leaf chlorophyll and photosynthetic capacity for improved modeling of GPP. *Agric. For. Meteorol.* 177, 10–23. doi:10.1016/j.agrformet.2013.04.006
- Hu, C., Chen, Z., Clayton, T.D., Swarzenski, P., Brock, J.C., Muller-Karger, F.E., 2004. Assessment of estuarine water-quality indicators using MODIS medium-resolution bands: Initial results from Tampa Bay, FL. *Remote Sens. Environ.* 93, 423–441. doi:10.1016/j.rse.2004.08.007
- Huang, C.-Y., Asner, G.P., 2009. Applications of Remote Sensing to Alien Invasive Plant Studies. *Sensors* 9, 4869–4889. doi:10.3390/s90604869
- Huang, S., Ding, J., Zou, J., Liu, B., Zhang, J., Chen, W., 2019. Soil moisture retrieval based on sentinel-1 imagery under sparse vegetation coverage. *Sensors* 19, 589. doi:10.3390/s19030589
- Hunt, E.R., Doraiswamy, P.C., McMurtrey, J.E., Daughtry, C.S.T., Perry, E.M., Akhmedov, B., 2013. A visible band index for remote sensing leaf chlorophyll content at the canopy scale. *Int. J. Appl. Earth Obs. Geoinf.* 21, 103–112. doi:10.1016/j.jag.2012.07.020
- Jackson, T.J., 2002. Remote sensing of soil moisture: Implications for groundwater recharge. *Hydrogeol. J.* 10, 40–51. doi:10.1007/s10040-001-0168-2
- Jackson, T.J., Chen, D., Cosh, M., Li, F., Anderson, M., Walthall, C., Doraiswamy, P., Hunt, E.R., 2004. Vegetation water content mapping using Landsat data derived normalized difference water index for corn and soybeans. *Remote Sens. Environ.* 92, 475–482. doi:10.1016/j.rse.2003.10.021
- Jain, M., Mondal, P., DeFries, R.S., Small, C., Galford, G.L., 2013. Mapping cropping intensity of smallholder farms: A comparison of methods using multiple sensors. *Remote Sens. Environ.* 134, 210–223. doi:10.1016/j.rse.2013.02.029
- Jakubauskas, M.E., Legates, D.R., Kastens, J.H., 2002. Crop identification using harmonic analysis of time-series AVHRR NDVI data. *Comput. Electron. Agric.* 37, 127–139. doi:10.1016/S0168-1699(02)00116-3
- Jennings, M.D., 2000. Gap analysis: Concepts, methods, and recent results. *Landsc. Ecol.* 15, 5–20. doi:10.1023/A:1008184408300
- Jiang, D., Wang, N., Yang, X., Wang, J., 2003. Study on the interaction between NDVI profile and the growing status of crops. *Chinese Geogr. Sci.* 13, 62–65. doi:10.1007/s11769-003-0086-4
- Johnson, D.M., 2013. A 2010 map estimate of annually tilled cropland within the conterminous United States. *Agric. Syst.* 114, 95–105. doi:10.1016/j.agsy.2012.08.004
- Johnston, M., Licker, R., Foley, J., Holloway, T., Mueller, N.D., Barford, C., Kucharik, C., 2011. Closing the gap: global potential for

- increasing biofuel production through agricultural intensification. *Environ. Res. Lett.* 6, 1–11. doi:10.1088/1748-9326/6/3/034028
- Kalubarme, M.H., Potdar, M.B., Manjunath, K.R., Mahey, R.K., Siddhu, S.S., 2003. Growth profile based crop yield models: A case study of large area wheat yield modelling and its extendibility using atmospheric corrected NOAA AVHRR data. *Int. J. Remote Sens.* 24, 2037–2054. doi:10.1080/01431160210156018
- Karnieli, A., Bayarjargal, Y., Bayasgalan, M., Mandakh, B., Dugarjav, C., Burgheimer, J., Khudulmur, S., Bazha, S.N., Gunin, P.D., 2013. Do vegetation indices provide a reliable indication of vegetation degradation? A case study in the Mongolian pastures. *Int. J. Remote Sens.* 34, 6243–6262. doi:10.1080/01431161.2013.793865
- Kasampalis, D., Alexandridis, T., Deva, C., Challinor, A., Moshou, D., Zalidis, G., 2018. Contribution of Remote Sensing on Crop Models: A Review. *J. Imaging* 4, 52. doi:10.3390/jimaging4040052
- Kasischke, E.S., Melack, J.M., Dobson, M.C., 1997. The use of imaging radars for ecological applications - A review. *Remote Sens. Environ.* 59, 141–156. doi:10.1016/S0034-4257(96)00148-4
- Kerr, J.T., Cihlar, J., 2003. Land use and cover with intensity of agriculture for Canada from satellite and census data. *Glob. Ecol. Biogeogr.* 12, 161–172. doi:10.1046/j.1466-822X.2003.00017.x
- Kerr, J.T., Ostrovsky, M., 2003. From space to species: Ecological applications for remote sensing. *Trends Ecol. Evol.* 18, 299–305. doi:10.1016/S0169-5347(03)00071-5
- Kerr, J.T., Southwood, T.R.E., Cihlar, J., 2001. Remotely sensed habitat diversity predicts butterfly species richness and community similarity in Canada. *PNAS* 98, 11365–11370. doi:10.1073/pnas.201398398
- Khan, N.M., Rastoskuev, V. V., Sato, Y., Shiozawa, S., 2005. Assessment of hydrosaline land degradation by using a simple approach of remote sensing indicators. *Agric. Water Manag.* 77, 96–109. doi:10.1016/j.agwat.2004.09.038
- Khanna, S., Palacios-Orueta, A., Whiting, M.L., Ustin, S.L., Riaño, D., Litago, J., 2007. Development of angle indexes for soil moisture estimation, dry matter detection and land-cover discrimination. *Remote Sens. Environ.* 109, 154–165. doi:10.1016/j.rse.2006.12.018
- Kimothi, M.M., Dasari, A., 2010. Methodology to map the spread of an invasive plant (*Lantana camara* L.) in forest ecosystems using Indian remote sensing satellite data. *Int. J. Remote Sens.* 31, 3273–3289. doi:10.1080/01431160903121126
- Kloiber, S.M., Brezonik, P.L., Olmanson, L.G., Bauer, M.E., 2002. A procedure for regional lake water clarity assessment using Landsat multispectral data. *Remote Sens. Environ.* 82, 38–47. doi:10.1016/S0034-4257(02)00022-6
- Koblinsky, C.J., Clarke, R.T., Brenner, A.C., Frey, H., 1993. Measurement of river level variations with satellite altimetry. *Water Resour. Res.* 29, 1839–1848. doi:10.1029/93WR00542
- Kuenzer, C., Ottinger, M., Wegmann, M., Guo, H., Wang, C., Zhang, J., Dech, S., Wikelski, M., 2014. Earth observation satellite sensors for biodiversity monitoring: potentials and bottlenecks. *Int. J. Remote Sens.* 35, 6599–6647. doi:10.1080/01431161.2014.964349
- Kumar, S., Gautam, G., Saha, S.K., 2015. Hyperspectral remote sensing data derived spectral indices in characterizing salt-affected soils: a case study of Indo-Gangetic plains of India. *Environ. Earth Sci.* 73, 3299–3308. doi:10.1007/s12665-014-3613-y
- Laba, M., Blair, B., Downs, R., Monger, B., Philpot, W., Smith, S., Sullivan, P., Baveye, P.C., 2010. Use of textural measurements to map invasive wetland plants in the Hudson River National Estuarine Research Reserve with IKONOS satellite imagery. *Remote Sens. Environ.* 114, 876–886. doi:10.1016/j.rse.2009.12.002
- Laba, M., Downs, R., Smith, S., Welsh, S., Neider, C., White, S., Richmond, M., Philpot, W., Baveye, P., 2008. Mapping invasive wetland plants in the Hudson River National Estuarine Research Reserve using quickbird satellite imagery. *Remote Sens. Environ.* 112, 286–300. doi:10.1016/j.rse.2007.05.003
- Lai, Y.R., Pringle, M.J., Kopittke, P.M., Menzies, N.W., Orton, T.G., Dang, Y.P., 2018. An empirical model for prediction of wheat yield, using time-integrated Landsat NDVI. *Int. J. Appl. Earth Obs. Geoinf.* 72, 99–108. doi:10.1016/j.jag.2018.07.013
- Lauver, C.L., 1997. Mapping species diversity patterns in the Kansas shortgrass region by integrating remote sensing and vegetation analysis. *J. Veg. Sci.* 8, 387–394. doi:10.2307/3237328

- Lavery, P., Pattiaratchi, C., Wyllie, A., Hick, P., 1993. Water Quality Monitoring in Estuarine Waters Using the Landsat Thematic Mapper. *Remote Sens. Environ.* 46, 268–280. doi:10.1016/0034-4257(93)90047-2
- Le Féon, V., Schermann-Legionnet, A., Delettre, Y., Aviron, S., Billeter, R., Bugter, R., Hendrickx, F., Burel, F., 2010. Intensification of agriculture, landscape composition and wild bee communities: A large scale study in four European countries. *Agric. Ecosyst. Environ.* 137, 143–150. doi:10.1016/j.agee.2010.01.015
- Leblon, B., Guerif, M., Baret, F., 1991. The use of remotely sensed data in estimation of PAR use efficiency and biomass production of flooded rice. *Remote Sens. Environ.* 38, 147–158. doi:10.1016/0034-4257(91)90076-I
- Li, L., Friedl, M.A., Xin, Q., Gray, J., Pan, Y., Froking, S., 2014. Mapping crop cycles in China using MODIS-EVI time series. *Remote Sens.* 6, 2473–2493. doi:10.3390/rs6032473
- Li, Z., Guo, X., 2012. Detecting climate effects on vegetation in northern mixed rairie using NOAA AVHRR 1-km time-series NDVI data. *Remote Sens.* 4, 120–134. doi:10.3390/rs4010120
- Li, Z., Xu, D., Guo, X., 2014. Remote sensing of ecosystem health: opportunities, challenges, and future perspectives. *Sensors* 14, 21117–21139. doi:10.3390/s141121117
- Liu, H., Li, Q., Shi, T., Hu, S., Wu, G., Zhou, Q., 2017. Application of Sentinel 2 MSI Images to Retrieve Suspended Particulate Matter Concentrations in Poyang Lake. *Remote Sens.* 9, 761. doi:10.3390/rs9070761
- Liu, J., Pattey, E., Miller, J.R., McNairn, H., Smith, A., Hu, B., 2010. Estimating crop stresses, aboveground dry biomass and yield of corn using multi-temporal optical data combined with a radiation use efficiency model. *Remote Sens. Environ.* 114, 1167–1177. doi:10.1016/j.rse.2010.01.004
- Lobell, D.B., Baldos, U.L.C., Hertel, T.W., 2013. Climate adaptation as mitigation: the case of agricultural investments. *Environ. Res. Lett.* 8, 1–12. doi:10.1088/1748-9326/8/1/015012
- Luoto, M., Kuussaari, M., Toivonen, T., 2002a. Modelling butterfly distribution based on remote sensing data. *J. Biogeogr.* 29, 1027–1037. doi:10.1046/j.1365-2699.2002.00728.x
- Luoto, M., Toivonen, T., Heikkinen, R.K., 2002b. Prediction of total and rare plant species richness in agricultural landscapes from satellite images and topographic data. *Landsc. Ecol.* 17, 195–217. doi:10.1023/A:1020288509837
- Mairota, P., Cafarelli, B., Didham, R.K., Lovergine, F.P., Lucas, R.M., Nagendra, H., Rocchini, D., Tarantino, C., 2015. Challenges and opportunities in harnessing satellite remote-sensing for biodiversity monitoring. *Ecol. Inform.* 30, 207–214. doi:10.1016/j.ecoinf.2015.08.006
- Martinez, J.M., Guyot, J.L., Cochonneau, G., 2008. Monitoring of surface water quality in large rivers with satellite imagery - Application to the Amazon basin. www.iwra.org/congress/resource/abs649_article.pdf.
- Mashimbye, Z.E., Cho, M.A., Nell, J.P., De clerck, W.P., Van niekerk, A., Turner, D.P., 2012. Model-Based Integrated Methods for Quantitative Estimation of Soil Salinity from Hyperspectral Remote Sensing Data: A Case Study of Selected South African Soils. *Pedosphere* 22, 640–649. doi:10.1016/S1002-0160(12)60049-6
- McGarigal, K., Cushman, S.A., Ene, E., 2012. FRAGSTATS v4: Spatial Pattern Analysis Program for Categorical Maps [WWW Document]. URL <http://www.umass.edu/landeco/research/fragstats/fragstats.html> (accessed 4.21.17).
- Metternicht, G.I., Zinck, J.A., 2003. Remote sensing of soil salinity: potentials and constraints. *Remote Sens. Environ.* 85, 1–20. doi:10.1016/S0034-4257(02)00188-8
- Metternicht, G.I., Zinck, J.A., 1998. Evaluating the information content of JERS-1 SAR and Landsat TM data for discrimination of soil erosion features. *ISPRS J. Photogramm. Remote Sens.* 53, 143–153. doi:10.1016/S0924-2716(98)00004-5
- Meynard, C.N., Quinn, J.F., 2007. Predicting species distributions: a critical comparison of the most common statistical models using artificial species. *J. Biogeogr.* 34, 1455–1469. doi:10.1111/j.1365-2699.2007.01720.x
- Michailovsky, C.I., McEnnis, S., Berry, P.A.M., Smith, R., Bauer-Gottwein, P., 2012. River monitoring from satellite radar altimetry in the Zambezi River basin. *Hydrol. Earth Syst. Sci.* 16, 2181–2192. doi:10.5194/hess-16-2181-2012
- Mingwei, Z., Qingbo, Z., Zhongxin, C., Jia, L., Yong, Z., Chongfa, C., 2008. Crop discrimination in Northern China with double cropping systems using Fourier analysis of time-series MODIS data. *Int. J. Appl. Earth Obs. Geoinf.* 10, 476–485. doi:10.1016/j.jag.2007.11.002

- Mirik, M., Ansley, R.J., 2012. Utility of Satellite and Aerial Images for Quantification of Canopy Cover and Infilling Rates of the Invasive Woody Species Honey Mesquite (*Prosopis Glandulosa*) on Rangeland. *Remote Sens.* 4, 1947–1962. doi:10.3390/rs4071947
- Monteith, J.L., 1972. Solar Radiation and Productivity in Tropical Ecosystems. *J. Appl. Ecol.* 9, 747–766. doi:10.2307/2401901
- Morel, J., Todoroff, P., Bégué, A., Bury, A., Martiné, J.F., Petit, M., 2014. Toward a satellite-based system of sugarcane yield estimation and forecasting in smallholder farming conditions: A case study on reunion island. *Remote Sens.* 6, 6620–6635. doi:10.3390/rs6076620
- Moriondo, M., Maselli, F., Bindi, M., 2007. A simple model of regional wheat yield based on NDVI data. *Eur. J. Agron.* 26, 266–274. doi:10.1016/j.eja.2006.10.007
- Mueller, N., Lewis, A., Roberts, D., Ring, S., Melrose, R., Sixsmith, J., Lymburner, L., McIntyre, A., Tan, P., Curnow, S., Ip, A., 2016. Water observations from space: Mapping surface water from 25years of Landsat imagery across Australia. *Remote Sens. Environ.* 174, 341–352. doi:10.1016/j.rse.2015.11.003
- Muñoz, J., Felicísimo, Á.M., 2004. Comparison of statistical methods commonly used in predictive modelling. *J. Veg. Sci.* 15, 285–292. doi:10.1111/j.1654-1103.2004.tb02263.x
- Mutiga, J.K., Su, Z., Woldai, T., 2010. Using satellite remote sensing to assess evapotranspiration: Case study of the upper Ewaso Ng'iro North Basin, Kenya. *Int. J. Appl. Earth Obs. Geoinf.* 12S, S100–S108. doi:10.1016/j.jag.2009.09.012
- Myneni, R.B., Nemani, R.R., Running, S.W., 1997. Estimation of global leaf area index and absorbed PAR using radiative transfer models. *IEEE Trans. Geosci. Remote Sens.* 35, 1380–1393. doi:10.1109/36.649788
- Nagendra, H., 2001. Using remote sensing to assess biodiversity. *Int. J. Remote Sens.* 22, 2377–2400. doi:10.1080/01431160117096
- Nagendra, H., Gadgil, M., 1999. Satellite Imagery as a Tool for Monitoring Species Diversity: An Assessment. *J. Appl. Ecol.* 36, 388–397. doi:10.1046/j.1365-2664.1999.00406.x
- Nagy, A., Fehér, J., Tamás, J., 2018. Wheat and maize yield forecasting for the Tisza river catchment using MODIS NDVI time series and reported crop statistics. *Comput. Electron. Agric.* 151, 41–49. doi:10.1016/j.compag.2018.05.035
- Ngoc, D.D., Loisel, H., Jamet, C., Vantrepotte, V., Duforêt-Gaurier, L., Minh, C.D., Mangin, A., 2019. Coastal and inland water pixels extraction algorithm (WiPE) from spectral shape analysis and HSV transformation applied to Landsat 8 OLI and Sentinel-2 MSI. *Remote Sens. Environ.* 223, 208–228. doi:10.1016/j.rse.2019.01.024
- Niedertscheider, M., Kastner, T., Fetzl, T., Haberl, H., Kroisleitner, C., Plutzer, C., Erb, K.-H., 2016. Mapping and analysing cropland use intensity from a NPP perspective. *Environ. Res. Lett.* 11. doi:10.1088/1748-9326/11/1/014008
- Numata, I., Roberts, D.A., Chadwick, O.A., Schimel, J., Sampaio, F.R., Leonidas, F.C., Soares, J. V., 2007. Characterization of pasture biophysical properties and the impact of grazing intensity using remotely sensed data. *Remote Sens. Environ.* 109, 314–327. doi:10.1016/j.rse.2007.01.013
- Olofsson, P., Lagergren, F., Lindroth, A., Lindström, J., Klemetsson, L., Kutsch, W., Eklundh, L., 2008. Towards operational remote sensing of forest carbon balance across Northern Europe. *Biogeosciences* 5, 817–832. doi:10.5194/bg-5-817-2008
- Pacheco, A., McNairn, H., 2010. Evaluating multispectral remote sensing and spectral unmixing analysis for crop residue mapping. *Remote Sens. Environ.* 114, 2219–2228. doi:10.1016/j.rse.2010.04.024
- Palacios-Orueta, A., Ustin, S.L., 1998. Remote sensing of soil properties in the Santa Monica Mountains I. Spectral analysis. *Remote Sens. Environ.* 65, 170–183. doi:10.1016/S0034-4257(98)00024-8
- Palmer, M.W., Earls, P.G., Hoagland, B.W., White, P.S., Wohlgemuth, T., 2002. Quantitative tools for perfecting species lists. *Environmetrics* 13, 121–137. doi:10.1002/env.516
- Pan, G., Sun, G.J., Li, F.M., 2009. Using QuickBird imagery and a production efficiency model to improve crop yield estimation in the semi-arid hilly Loess Plateau, China. *Environ. Model. Softw.* 24, 510–516. doi:10.1016/j.envsoft.2008.09.014
- Patel, N.R., Bhattacharjee, B., Mohammed, A.J., Tanupriya, B., Saha, S.K., 2006. Remote sensing of regional yield assessment of wheat in Haryana, India. *Int. J. Remote Sens.* 27, 4071–4090. doi:10.1080/01431160500377188

- Pereira-Sandoval, M., Ruiz-Verdu, A., Tenjo, C., Delegido, J., Urrego, P., Pena, R., Vicente, E., Soria, J., Soria, J., Moreno, J., 2018. Calibration and Validation of Algorithms for the Estimation of Chlorophyll-A Concentration and Secchi Depth in Inland Waters with Sentinel-2. *Limnetica* 38, 471–487. doi:10.1109/igarss.2018.8517371
- Petropoulos, G.P., Srivastava, P.K., Piles, M., Pearson, S., 2018. Earth observation-based operational estimation of soil moisture and evapotranspiration for agricultural crops in support of sustainable water management. *Sustainability* 10, 181. doi:10.3390/su10010181
- Ponnuram, G.G., Rao, Y.S., 2018. The application of compact polarimetric decomposition algorithms to L-band PolSAR data in agricultural areas. *Int. J. Remote Sens.* 39, 8337–8360. doi:10.1080/01431161.2018.1488281
- Potter, P., Ramankutty, N., Bennett, E.M., Donner, S.D., 2010. Characterizing the spatial patterns of global fertilizer application and manure production. *Earth Interact.* 14, 1–22. doi:10.1175/2009EI288.1
- Prasad, A.K., Chai, L., Singh, R.P., Kafatos, M., 2006. Crop yield estimation model for Iowa using remote sensing and surface parameters. *Int. J. Appl. Earth Obs. Geoinf.* 8, 26–33. doi:10.1016/j.jag.2005.06.002
- Qian, T., Tsunekawa, A., Peng, F., Masunaga, T., Wang, T., Li, R., 2019. Derivation of salt content in salinized soil from hyperspectral reflectance data: A case study at Minqin Oasis, Northwest China. *J. Arid Land* 11, 111–122. doi:10.1007/s40333-019-0091-9
- Rahman, M.M., Moran, M.S., Thoma, D.P., Bryant, R., Holifield Collins, C.D., Jackson, T., Orr, B.J., Tischler, M., 2008. Mapping surface roughness and soil moisture using multi-angle radar imagery without ancillary data. *Remote Sens. Environ.* 112, 391–402. doi:10.1016/j.rse.2006.10.026
- Ravan, S.A., Roy, P.S., Sharma, C.M., 1995. Space remote sensing for spatial vegetation characterization. *J. Biosci* 20, 427–438.
- Reidsma, P., Tekelenburg, T., Van Den Berg, M., Alkemade, R., 2006. Impacts of land-use change on biodiversity: An assessment of agricultural biodiversity in the European Union. *Agric. Ecosyst. Environ.* 114, 86–102. doi:10.1016/j.agee.2005.11.026
- Ren, J., Chen, Z., Zhou, Q., Tang, H., 2008. Regional yield estimation for winter wheat with MODIS-NDVI data in Shandong, China. *Int. J. Appl. Earth Obs. Geoinf.* 10, 403–413. doi:10.1016/j.jag.2007.11.003
- Ritchie, J.C., Cooper, C.M., 2001. Remote sensing techniques for determining water quality: Applications to TMDLs, in: TMDL Science Issues Conference (2001), Water Environment Federation, Alexandria, VA. pp. 367–374.
- Rocchini, D., Balkenhol, N., Carter, G.A., Foody, G.M., Gillespie, T.W., He, K.S., Kark, S., Levin, N., Lucas, K., Luoto, M., Nagendra, H., Oldeland, J., Ricotta, C., Southworth, J., Neteler, M., 2010. Remotely sensed spectral heterogeneity as a proxy of species diversity: Recent advances and open challenges. *Ecol. Inform.* 5, 318–329. doi:10.1016/j.ecoinf.2010.06.001
- Rocchini, D., Ricotta, C., Chiarucci, A., 2007. Using satellite imagery to assess plant species richness: The role of multispectral systems. *Appl. Veg. Sci.* 10, 325–331. doi:10.1111/j.1654-109X.2007.tb00431.x
- Ruimy, A., Saugier, B., 1994. Methodology for the estimation of terrestrial net primary production from remotely sensed data. *J. Geophys. Res.* 99, 5263–5283. doi:10.1029/93JD03221
- Sakowska, K., Juszczak, R., Gianelle, D., 2016. Remote Sensing of Grassland Biophysical Parameters in the Context of the Sentinel-2 Satellite Mission. *J. Sensors* 2016, 1–16. doi:10.1155/2016/4612809
- Salazar, L., Kogan, F., Roytman, L., 2007. Use of remote sensing data for estimation of winter wheat yield in the United States. *Int. J. Remote Sens.* 28, 3795–3811. doi:10.1080/01431160601050395
- Saura, S., 2004. Effects of remote sensor spatial resolution and data aggregation on selected fragmentation indices. *Landsc. Ecol.* 19, 197–209. doi:10.1023/B:LAND.0000021724.60785.65
- Saveraid, E.H., Debinski, D.M., Kindscher, K., Jakubauskas, M.E., 2001. A comparison of satellite data and landscape variables in predicting bird species occurrences in the Greater Yellowstone Ecosystem, USA. *Landsc. Ecol.* 16, 71–83. doi:10.1023/A:1008119219788
- Scharf, P., Schmidt, J.P., Kitchen, N.R., Sudduth, K.A., 2002. Remote sensing for nitrogen management. *J. Soil Water Conserv.* 57, 518–524.
- Schino, G., Borfecchia, F., De Cecco, L., Dibari, C., Iannetta, M., Martini, S., Pedrotti, F., 2003. Satellite estimate of grass biomass in a mountainous range in central Italy. *Agrofor. Syst.* 59, 157–162. doi:10.1023/A:1026308928874

- Schumaker, N.H., 1996. Using landscape indices to predict habitat connectivity. *Ecology* 77, 1210–1225. doi:10.2307/2265590
- Sharma, K.D., Singh, S., 1995. Satellite remote sensing for soil erosion modelling using the ANSWERS model. *Hydrol. Sci.* 40, 259–272. doi:10.1080/02626669509491408
- Shoshany, M., Goldshleger, N., Chudnovsky, A., 2013. Monitoring of agricultural soil degradation by remote-sensing methods: a review. *Int. J. Remote Sens.* 34, 6152–6181. doi:10.1080/01431161.2013.793872
- Shuai, G., Zhang, J., Basso, B., Pan, Y., Zhu, X., Zhu, S., Liu, H., 2019. Multi-temporal RADARSAT-2 polarimetric SAR for maize mapping supported by segmentations from high-resolution optical image. *Int. J. Appl. Earth Obs. Geoinf.* 74, 1–15. doi:10.1016/j.jag.2018.08.021
- Sibanda, M., Mutanga, O., Rouget, M., 2015. Examining the potential of Sentinel-2 MSI spectral resolution in quantifying above ground biomass across different fertilizer treatments. *ISPRS J. Photogramm. Remote Sens.* 110, 55–65. doi:10.1016/j.isprsjprs.2015.10.005
- Siebert, S., Portmann, F.T., Döll, P., 2010. Global patterns of cropland use intensity. *Remote Sens.* 2, 1625–1643. doi:10.3390/rs2071625
- Singh, R., Semwal, D.P., Rai, A., Chhikara, R.S., 2002. Small area estimation of crop yield using remote sensing satellite data. *Int. J. Remote Sens.* 23, 49–56. doi:10.1080/01431160010014756
- Smith, L.C., 1997. Satellite remote sensing of river inundation area, stage, and discharge: a review. *Hydrol. Process.* 11, 1427–1439. doi:10.1002/(SICI)1099-1085(199708)11:10<1427::AID-HYP473>3.0.CO;2-S
- Son, N.T., Chen, C.F., Chen, C.R., Minh, V.Q., Trung, N.H., 2014. A comparative analysis of multitemporal MODIS EVI and NDVI data for large-scale rice yield estimation. *Agric. For. Meteorol.* 197, 52–64. doi:10.1016/j.agrformet.2014.06.007
- Sòria-perpinyà, X., Urrego, P., Pereira-sandoval, M., Ruiz-verdú, A., Peña, R., Soria, J.M., Delegido, J., Vicente, E., Moreno, J., València, U. De, José, C.C., Martínez, B., Martínez, B., 2019. Monitoring the ecological state of a hypertrophic lake (Albufera of València, Spain) using multitemporal Sentinel-2 images 38, 457–469. doi:10.23818/limn.38.26
- Srivastava, P.K., 2017. Satellite Soil Moisture: Review of Theory and Applications in Water Resources. *Water Resour. Manag.* 31, 3161–3176. doi:10.1007/s11269-017-1722-6
- Stevens, A., Udelhoven, T., Denis, A., Tychon, B., Liou, R., Hoffmann, L., van Wesemael, B., 2010. Measuring soil organic carbon in croplands at regional scale using airborne imaging spectroscopy. *Geoderma* 158, 32–45. doi:10.1016/j.geoderma.2009.11.032
- Stickler, C.M., Southworth, J., 2008. Application of multi-scale spatial and spectral analysis for predicting primate occurrence and habitat associations in Kibale National Park, Uganda. *Remote Sens. Environ.* 112, 2170–2186. doi:10.1016/j.rse.2007.10.013
- Sulistioadi, Y.B., Tseng, K.H., Shum, C.K., Hidayat, H., Sumaryono, M., Suhardiman, A., Setiawan, F., Sunarso, S., 2015. Satellite radar altimetry for monitoring small rivers and lakes in Indonesia. *Hydrol. Earth Syst. Sci.* 19, 341–359. doi:10.5194/hess-19-341-2015
- Taghadosi, M.M., Hasanlou, M., Eftekhari, K., 2019. Soil salinity mapping using dual-polarized SAR Sentinel-1 imagery. *Int. J. Remote Sens.* 40, 237–252. doi:10.1080/01431161.2018.1512767
- Taylor, G.R., Mah, A.H., Kruse, F.A., Kierein-Young, K.S., Hewson, R.D., Bennett, B.A., 1996. Characterization of Saline Soils Using Airborne Radar Imagery. *Remote Sens. Environ.* 57, 127–142. doi:10.1016/0034-4257(95)00239-1
- Teillard, F., Allaire, G., Cahuzac, E., Léger, F., Maigné, E., Tichit, M., 2012. A novel method for mapping agricultural intensity reveals its spatial aggregation: Implications for conservation policies. *Agric. Ecosyst. Environ.* 149, 135–143. doi:10.1016/j.agee.2011.12.018
- Temme, A.J.A.M., Verburg, P.H., 2011. Mapping and modelling of changes in agricultural intensity in Europe. *Agric. Ecosyst. Environ.* 140, 46–56. doi:10.1016/j.agee.2010.11.010
- Thenkabail, P.S., Biradar, C.M., Noojipady, P., Dheeravath, V., Li, Y., Velpuri, M., Gumma, M., Gangalakunta, O.R.P., Turral, H., Cai, X., Vithanage, J., Schull, M. a., Dutta, R., 2009. Global irrigated area map (GIAM), derived from remote sensing, for the end of the last millennium. *Int. J. Remote Sens.* 30, 3679–3733. doi:10.1080/01431160802698919

- Tilling, A.K., O'Leary, G.J., Ferwerda, J.G., Jones, S.D., Fitzgerald, G.J., Rodriguez, D., Belford, R., 2007. Remote sensing of nitrogen and water stress in wheat. *F. Crop. Res.* 104, 77–85. doi:10.1016/j.fcr.2007.03.023
- Trivero, P., Borasi, M., Biamino, W., Cavagnero, M., Rinaudo, C., Bonansea, M., Lanfri, S., 2013. River pollution remediation monitored by optical and infrared high-resolution satellite images. *Environ. Monit. Assess.* 185, 7647–7658. doi:10.1007/s10661-013-3125-3
- Trombetti, M., Riaño, D., Rubio, M.A., Cheng, Y.B., Ustin, S.L., 2008. Multi-temporal vegetation canopy water content retrieval and interpretation using artificial neural networks for the continental USA. *Remote Sens. Environ.* 112, 203–215. doi:10.1016/j.rse.2007.04.013
- Tsai, F., Chou, M.J., 2006. Texture augmented analysis of high resolution satellite imagery in detecting invasive plant species. *J. Chinese Inst. Eng.* 29, 581–592. doi:10.1080/02533839.2006.9671155
- Tsai, F., Lin, E.K., Yoshino, K., 2007. Spectrally segmented principal component analysis of hyperspectral imagery for mapping invasive plant species. *Int. J. Remote Sens.* 28, 1023–1039. doi:10.1080/01431160600887706
- Tulbure, M.G., Broich, M., 2013. Spatiotemporal dynamic of surface water bodies using Landsat time-series data from 1999 to 2011. *ISPRS J. Photogramm. Remote Sens.* 79, 44–52. doi:10.1016/j.isprsjprs.2013.01.010
- Turner, D.P., Cohen, W.B., Kennedy, R.E., Fassnacht, K.S., Briggs, J.M., 1999. Relationships between Leaf Area Index and Landsat TM Spectral Vegetation Indices across Three Temperate Zone Sites. *Remote Sens. Environ.* 70, 52–68. doi:10.1016/S0034-4257(99)00057-7
- Turner, W., Spector, S., Gardiner, N., Fladeland, M., Sterling, E., Steininger, M., 2003. Remote sensing for biodiversity science and conservation. *Trends Ecol. Evol.* 18, 306–314. doi:10.1016/S0169-5347(03)00070-3
- van Leeuwen, M., Nieuwenhuis, M., 2010. Retrieval of forest structural parameters using LiDAR remote sensing. *Eur. J. For. Res.* 129, 749–770. doi:10.1007/s10342-010-0381-4
- Vannier, C., Vasseur, C., Hubert-Moy, L., Baudry, J., 2011. Multiscale ecological assessment of remote sensing images. *Landsc. Ecol.* 26, 1053–1069. doi:10.1007/s10980-011-9626-y
- Verhoef, W., 1984. Light scattering by leaf layers with application to canopy reflectance modeling: The SAIL model. *Remote Sens. Environ.* 16, 125–141. doi:10.1016/0034-4257(84)90057-9
- Verpoorter, C., Kutser, T., Tranvik, L., 2012. Automated mapping of water bodies using Landsat multispectral data. *Limnol. Oceanogr. Methods* 10, 1037–1050. doi:10.4319/lom.2012.10.1037
- Verstraeten, W.W., Veroustraete, F., Van Der Sande, C.J., Grootaers, I., Feyen, J., 2006. Soil moisture retrieval using thermal inertia, determined with visible and thermal spaceborne data, validated for European forests. *Remote Sens. Environ.* 101, 299–314. doi:10.1016/j.rse.2005.12.016
- Vibhute, A.D., Gawali, B.W., 2013. Analysis and Modeling of Agricultural Land use using Remote Sensing and Geographic Information System: a Review. *Int. J. Eng. Res. Appl.* 3, 81–91.
- Vignolo, A., Pochettino, A., Cicerone, D., 2006. Water quality assessment using remote sensing techniques: Medrano Creek, Argentina. *J. Environ. Manage.* 81, 429–433. doi:10.1016/j.jenvman.2005.11.019
- Vrieling, A., 2006. Satellite remote sensing for water erosion assessment: A review. *Catena* 65, 2–18. doi:10.1016/j.catena.2005.10.005
- Wagner, W., Naeimi, V., Scipal, K., Jeu, R., Martínez-Fernández, J., 2007. Soil moisture from operational meteorological satellites. *Hydrogeol. J.* 15, 121–131. doi:10.1007/s10040-006-0104-6
- Walsh, S.J., McCleary, A.L., Mena, C.F., Shao, Y., Tuttle, J.P., González, A., Atkinson, R., 2008. QuickBird and Hyperion data analysis of an invasive plant species in the Galapagos Islands of Ecuador: Implications for control and land use management. *Remote Sens. Environ.* 112, 1927–1941. doi:10.1016/j.rse.2007.06.028
- Wang, B., Jia, K., Liang, S., Xie, X., Wei, X., Zhao, X., Yao, Y., Zhang, X., 2018. Assessment of Sentinel-2 MSI Spectral Band Reflectances for Estimating Fractional Vegetation Cover. *Remote Sens.* 10, 1927. doi:10.3390/rs10121927
- Wang, K., Franklin, S.E., Guo, X., Cattet, M., 2010. Remote sensing of ecology, biodiversity and conservation: A review from the perspective of remote sensing specialists. *Sensors* 10, 9647–9667. doi:10.3390/s101109647

- Wang, L., Huang, J., Du, Y., Hu, Y., Han, P., 2013. Dynamic assessment of soil erosion risk using landsat TM and HJ satellite data in danjiangkou reservoir area, China. *Remote Sens.* 5, 3826–3848. doi:10.3390/rs5083826
- Wang, L., Qu, J.J., 2009. Satellite remote sensing applications for surface soil moisture monitoring: A review. *Front. Earth Sci. China* 3, 237–247. doi:10.1007/s11707-009-0023-7
- Wardlow, B.D., Egbert, S.L., 2008. Large-area crop mapping using time-series MODIS 250 m NDVI data: An assessment for the U.S. Central Great Plains. *Remote Sens. Environ.* 112, 1096–1116. doi:10.1016/j.rse.2007.07.019
- Wardlow, B.D., Egbert, S.L., Kastens, J.H., 2007. Analysis of time-series MODIS 250m vegetation index data for crop classification in the U.S. Central Great Plains. *Remote Sens. Environ.* 108, 290–310. doi:10.1016/j.rse.2006.11.021
- Weiers, S., Bock, M., Wissen, M., Rossner, G., 2004. Mapping and indicator approaches for the assessment of habitats at different scales using remote sensing and GIS methods. *Landsc. Urban Plan.* 67, 43–65. doi:10.1016/S0169-2046(03)00028-8
- Wen, X., Yang, X., 2010. Monitoring of water quality using remote sensing techniques. *Appl. Mech. Mater.* 29–32, 2360–2364. doi:10.4028/www.scientific.net/AMM.29-32.2360
- Weng, Y., Gong, P., Zhu, Z., 2008. Soil salt content estimation in the Yellow River delta with satellite hyperspectral data. *Can. J. Remote Sens.* 34, 259–270. doi:10.5589/m08-017
- Wolter, P.T., Johnston, C.A., Niemi, G.J., 2005. Mapping submergent aquatic vegetation in the US Great Lakes using Quickbird satellite data. *Int. J. Remote Sens.* 26, 5255–5274. doi:10.1080/01431160500219208
- Wu, B., Gommers, R., Zhang, M., Zeng, H., Yan, N., Zou, W., Zheng, Y., Zhang, N., Chang, S., Xing, Q., van Heijden, A., 2015. Global crop monitoring: A satellite-based hierarchical approach. *Remote Sens.* 7, 3907–3933. doi:10.3390/rs70403907
- Wu, B., Meng, J., Li, Q., Yan, N., Du, X., Zhang, M., 2014. Remote sensing-based global crop monitoring: Experiences with China's CropWatch system. *Int. J. Digit. Earth* 7, 113–137. doi:10.1080/17538947.2013.821185
- Wu, C., Wu, J., Qi, J., Zhang, L., Huang, H., Lou, L., Chen, Y., 2010. Empirical estimation of total phosphorus concentration in the mainstream of the Qiantang River in China using Landsat TM data. *Int. J. Remote Sens.* 31, 2309–2324. doi:10.1080/01431160902973873
- Wulder, M.A., Hall, R.J., Coops, Nicholas, C., Franklin, S.E., 2004. High spatial resolution remotely sensed data for ecosystem characterisation. *Bioscience* 54, 511–521.
- Xiao, X., Jian, X., Xiongfei, W., Chengfang, H., Xuejun, C., Zhaohui, W., Dengzhong, Z., 2015. Evaluation method of water quality for river based on multi-spectral remote sensing data, in: *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences - ISPRS Archives*. doi:10.5194/isprsarchives-XL-7-W3-1517-2015
- Xie, H., Zou, J., Jiang, H., Zhang, N., Choi, Y., 2014. Spatiotemporal pattern and driving forces of arable land-use intensity in China: Toward sustainable land management using emergy analysis. *Sustainability* 6, 3504–3520. doi:10.3390/su6063504
- Xu, C., Li, Y., Hu, J., Yang, X., Sheng, S., Liu, M., 2012. Evaluating the difference between the normalized difference vegetation index and net primary productivity as the indicators of vegetation vigor assessment at landscape scale. *Environ. Monit. Assess.* 184, 1275–1286. doi:10.1007/s10661-011-2039-1
- Xu, L., Zhang, H., Wang, C., Zhang, B., Liu, M., 2019. Crop classification based on temporal information using Sentinel-1 SAR time-series data. *Remote Sens.* 11, 53. doi:10.3390/rs11010053
- Yan, H., Xiao, X., Huang, H., Liu, J., Chen, J., Bai, X., 2014. Multiple Cropping Intensity in China Derived from Agro-meteorological Observations and MODIS Data. *Chinese Geogr. Sci.* 24, 205–219. doi:10.1007/s11769-013-0637-2
- Yang, C., Everitt, J.H., Bradford, J.M., 2006. Comparison of QuickBird satellite imagery and airborne imagery for mapping grain sorghum yield patterns. *Precis. Agric.* 7, 33–44. doi:10.1007/s11119-005-6788-0
- Yang, X., 2005. Remote sensing and GIS applications for estuarine ecosystem analysis: An overview. *Int. J. Remote Sens.* 26, 5347–5356. doi:10.1080/01431160500219406
- Zhang, M., Wu, B., Yu, M., Zou, W., Zheng, Y., 2014. Crop condition assessment with adjusted NDVI using the uncropped arable land ratio. *Remote Sens.* 6, 5774–5794. doi:10.3390/rs6065774

- Zhang, W., Li, H., 2016. Characterizing and Assessing the Agricultural Land Use Intensity of the Beijing Mountainous Region. *Sustainability* 8, 1–18. doi:10.3390/su8111180
- Zhang, Y., Atkinson, P.M., Li, X., Ling, F., Wang, Q., Du, Y., 2017. Learning-Based Spatial-Temporal Superresolution Mapping of Forest Cover with MODIS Images. *IEEE Trans. Geosci. Remote Sens.* 55, 600–614. doi:10.1109/TGRS.2016.2613140